

GLOBAL PROPERTIES OF TIGHT REEB FLOWS WITH APPLICATIONS TO FINSLER GEODESIC FLOWS ON S^2

UMBERTO L. HRYNIEWICZ AND PEDRO A. S. SALOMÃO

ABSTRACT. We show that if a Finsler metric on S^2 with reversibility r has flag curvatures K satisfying $(\frac{r}{r+1})^2 < K \leq 1$, then closed geodesics with specific contact-topological properties cannot exist, in particular there are no closed geodesics with precisely one transverse self-intersection point. This is a special case of a more general phenomenon, and other closed geodesics with many self-intersections are also excluded. We provide examples of Randers type, obtained by suitably modifying the metrics constructed by Katok [21], proving that this pinching condition is sharp. Our methods are borrowed from the theory of pseudo-holomorphic curves in symplectizations. Finally, we study global dynamical aspects of 3-dimensional energy levels C^2 -close to S^3 .

1. INTRODUCTION AND MAIN RESULTS

Classical and recent results show that pinching conditions on the curvatures of a Riemannian metric force the geodesic flow to present specific global behavior, usually encoded in geometric-topological and dynamical properties of closed geodesics. The interest in such phenomena can be traced back to Poincaré [23] and Birkhoff [9] where, among many other topics, the geodesic flow on positively curved surfaces was studied.

In the 1970s and 1980s this subject again received much attention. For example, in the work of Thorbergsson [27], Ballmann, Thorbergsson and Ziller [5, 6, 7] and Klingenberg [22] one finds many results relating pinching conditions on the curvatures to the existence (or non-existence) of closed geodesics with various topological and dynamical properties. Let us recall two theorems along these lines proved around the same time. As usual, a Riemannian metric is called δ -pinched if all sectional curvatures K satisfy $\delta \leq K \leq 1$.

Theorem 1.1 (Ballmann [4]). *Given $k \geq 1$ and $\epsilon > 0$, there exists $\delta < 1$ such that every prime closed geodesic of a δ -pinched metric on S^2 is either a simple curve of length in $[2\pi - \epsilon, 2\pi + \epsilon]$, or has at least k self-intersections and length $> \epsilon^{-1}$.*

Theorem 1.2 (Bangert [3]). *For every $\epsilon > 0$ there exists $\delta < 1$ such that the length l of every prime closed geodesic of a δ -pinched metric on S^n satisfies either $l \in [2\pi - \epsilon, 2\pi + \epsilon]$ or $l > \epsilon^{-1}$.*

Both theorems are of a perturbative nature and exhibit a “short-long” dichotomy for prime closed geodesics: if the metric is sufficiently pinched then their lengths are either close to the lengths in the round case (short), or arbitrarily large. This is surprising since one could try to imagine a sequence of metrics converging in C^2 to the round sphere admitting prime closed geodesics with lengths close to $2k\pi$, for some $k \geq 2$. However, this does not happen.

Motivated by the above statements one might consider the following questions in the more general framework of Finsler metrics on S^2 , or even in broader classes of Hamiltonian systems.

- a) How much can we relax the pinching of the flag curvatures of a (possibly non-reversible) Finsler metric on the 2-sphere and still keep some kind of dichotomy similar to that in Theorem 1.1?
- b) Can the “short-long” length dichotomy in Theorem 1.2 be generalized to a “low-high” action dichotomy on a broader class of Hamiltonian systems? If so, is there any additional topological information that can be extracted in low dimensions, in a way similar to Theorem 1.1?

Ballmann, Thorbergsson and Ziller [7] observe that a δ -pinched Riemannian metric on the 2-sphere, with $\delta > 1/4$, does not admit a closed geodesic with precisely one self-intersection. The proof is an immediate application of two well-known comparison theorems. To be more precise, the pinching condition $0 < \delta < K \leq 1$ implies Klingenberg’s estimate for the injectivity radius $\text{inj}(p) \geq \pi, \forall p \in S^2$. Since a closed geodesic γ with exactly one self-intersection point is the union of two loops, its length l must satisfy $l \geq 4\pi$. On the other hand, since γ is also a convex geodesic polygon, we have from Toponogov’s theorem the estimate $l \leq 2\pi/\sqrt{\delta} < 4\pi$ if $\delta > 1/4$. These two inequalities on the length l imply that such closed geodesic cannot exist. In this case, closed geodesics are either simple with length $\leq 2\pi/\sqrt{\delta}$, or have at least two self-intersections and length $\geq 6\pi$. This may be thought of as a simplest answer to a) in the Riemannian case, but maybe other pinching conditions will rule out other types of geodesics.


We use the theory of pseudo-holomorphic curves in symplectizations developed by H. Hofer, K. Wysocki and E. Zehnder as an alternative to comparison theorems. These methods reveal a more general phenomenon, in fact, under a certain pinching condition on the flag curvatures (cf. Theorem 1.5) there exists a larger class of immersed curves that cannot be realized as closed geodesics. This class includes curves with precisely one transverse self-intersection, but also many other curves with an arbitrarily large number of self-intersections. Then we exhibit examples of Randers type showing that the above mentioned pinching condition is optimal.

Finally, we quickly address b). It follows trivially from the method of Bangert [3] that a low-high action dichotomy holds for convex energy levels in $\mathbb{R}^{2n}$ which are C^2 -close to S^{2n-1} . In the case $n = 2$ we study the linking number between high- and low-action orbits.

1.1. Main results. We consider a weakened version of the notion of flat knot types discussed in [1], which relates to V. I. Arnold’s J^+ -theory of plane curves described in [2].

Definition 1.3. Consider the set $\mathcal{B}$ of C^2 -immersions $\gamma : S^1 \rightarrow S^2$ such that all self-intersections are either transverse or negative tangencies, i.e., if $\gamma(t_0) = \gamma(t_1)$ and $t_0 \neq t_1$ then $\dot{\gamma}(t_1) \notin \mathbb{R}^+ \dot{\gamma}(t_0)$. We say two curves $\gamma_0, \gamma_1 \in \mathcal{B}$ are equivalent if they are homotopic through curves in $\mathcal{B}$. A weak flat knot type is an equivalence class of curves in $\mathcal{B}$.

If we fix a Finsler metric F on S^2 then the unit sphere bundle $SS^2 := \{v \in TS^2 \mid F(v) = 1\}$ admits a contact form α_F given by the pull-back of the tautological 1-form of T^*S^2 via the associated Legendre transform. Any weak flat knot type of some $\gamma \in \mathcal{B}$ singles out a transverse knot type in the contact manifold $(SS^2, \ker \alpha_F)$

FIGURE 1. The weak flat knot type k_8 .

determined by the knot $\dot{\gamma}/F(\dot{\gamma})$. In particular, topological and contact invariants of transverse knots, like the self-linking number (cf. § 2.2 below), induce invariants of weak flat knot types on S^2 . Note that any prime closed geodesic of a Finsler metric on S^2 represents a weak flat knot type¹.

To give a computable concrete example, consider the weak flat knot type k_8 of an “eight-like curve” having precisely one self-intersection point which is transverse. The proof of the following lemma is found in § 3.1.

Lemma 1.4. *Let the C^2 -immersion $c : S^1 \rightarrow S^2$ represent the weak flat knot type k_8 , and let F be any Finsler metric on S^2 . Then the curve $\dot{c}/F(\dot{c})$ in the unit sphere bundle is unknotted and has self-linking number -1 .*

Before stating our main result we need to recall the notion of reversibility of a Finsler metric F , defined by Rademacher [24] as

$$(1) \quad r := \max\{F(-v) \mid F(v) = 1\} \geq 1.$$

It equals 1 exactly when $F(v) = F(-v) \forall v$, and F is called reversible in this case. The notion of reversibility is an essential ingredient in Rademacher’s proof of his sphere theorem for Finsler metrics.

Theorem 1.5. *The following assertions hold.*

- (i) *Let F be a Finsler metric on S^2 with reversibility r . If all flag curvatures K satisfy*

$$(2) \quad \left(\frac{r}{1+r} \right)^2 < K \leq 1$$

then no prime closed geodesic γ represents the weak flat knot type k_8 .

- (ii) *Statement (i) is optimal in the following sense: for every choice of $r \geq 1$ and $0 < \delta < (r/(r+1))^2$ there exists a Finsler metric on S^2 with reversibility r and δ -pinched flag curvatures admitting closed geodesics with precisely one transverse self-intersection.*

We stress the fact that the proof of part (i) in Theorem 1.5 does not make use of any version of Toponogov’s theorem for Finsler geometry. In fact, our method seems to be an alternative tool in those cases where such a comparison theorem may not be effective.

Note that there exist immersions $\gamma \in \mathcal{B}$ representing k_8 with an arbitrarily large number of self-intersections, see Figure 1 for an example with 3 self-intersections.

¹In the reversible case negative self-tangencies of closed prime geodesics never happen, so one gets a so-called flat knot type as defined in [1]. However, negative self-tangencies could appear in the non-reversible case.

Such immersions cannot be realized as a closed geodesic under the pinching condition (2).

To prove assertion (ii) we modify the metrics of Katok [21]. The examples are Randers metrics given by suitably chosen Zermelo navigation data on surfaces of revolution in $\mathbb{R}^3$, see § 3.3 for the detailed construction. Assertion (i) is proved by an application of pseudo-holomorphic curve theory in symplectizations, as introduced by Hofer in [14], developed by Hofer, Wysocki and Zehnder during the 1990s, and later by many other authors. The arguments are based on a dynamical characterization of the tight 3-sphere from [20], extending earlier results from [16, 17]. For an outline of the proof we refer to § 1.2.1 below.

Our second result relates to question b). Denote by $\mathbf{Conv}(2m)$ the set of closed and strictly convex hypersurfaces of class C^2 in $\mathbb{R}^{2m}$, equipped with the C^2 -topology. $\mathbb{R}^{2m}$ is endowed with its standard symplectic structure ω_0 , and each $S \in \mathbf{Conv}(2m)$ is oriented as the boundary of the bounded connected component of $\mathbb{R}^{2m} \setminus S$. A closed characteristic on S is a closed leaf of $TS^\perp \subset TS$, where $\perp$ denotes the ω_0 -symplectic orthogonal. These are precisely geometric images of closed Hamiltonian orbits, for any Hamiltonian realizing S as a regular energy level. We denote by $\mathcal{P}(S)$ the set of closed characteristics and think its elements as first iterates of periodic orbits of a Hamiltonian system. The action of a given $P \in \mathcal{P}(S)$ is $\mathcal{A}(P) = \int_P \lambda_0$, where $\lambda_0 = \frac{1}{2} \sum qdp - pdq$ is the standard Liouville form.

These Hamiltonian systems generalize geodesic flows on S^2 . In fact, as is well known, the geodesic flow of any Finsler metric on S^2 lifts to a Hamiltonian flow on a suitable star-shaped hypersurface in $\mathbb{R}^4$ via a double cover. This lifting procedure is nicely described in [13]. In general, however, S is not convex. If the metric is C^2 -close to the round metric then S belongs to $\mathbf{Conv}(4)$ and is close to S^3 . Action of an orbit on S coincides with length of its projection on S^2 . With this picture in mind we make the following statement.

Theorem 1.6. *Given $\epsilon > 0$ there exists a neighborhood $\mathcal{U}_\epsilon$ of S^{2m-1} in $\mathbf{Conv}(2m)$ such that if $S \in \mathcal{U}_\epsilon$ then every $P \in \mathcal{P}(S)$ satisfies $\mathcal{A}(P) \in [\pi - \epsilon, \pi + \epsilon]$ (short orbits) or $\mathcal{A}(P) > \epsilon^{-1}$ (long orbits). In the case $m = 2$, given any $k \geq 1$ there exists a neighborhood $\mathcal{U}_{\epsilon,k} \subset \mathcal{U}_\epsilon$ of S^3 such that if $S \in \mathcal{U}_{\epsilon,k}$ then $\text{link}(P, P') \geq k$ whenever P is short and P' is long.*

The assertion about the high-low action dichotomy is a direct application of Proposition 4.1 due to Bangert [3], and in the case $m = 2$ it is crucial to estimating the linking numbers. Obviously the short orbits are unknotted, have self-linking number -1 and their Conley-Zehnder indices belong to $\{3, 4, 5\}$. An analogous statement is true for energy levels C^2 -close to irrational ellipsoids, except that the high-low action dichotomy is trivial in this case. For an idea of the proof see § 1.2.2 below.

1.2. Outline of the main arguments. For convenience of the reader we sketch some of the main steps in the proofs of our results.

1.2.1. Non-existence of geodesics. Here we briefly explain why assertion (i) of Theorem 1.5 holds. For more details see § 3.2.

A contact form λ on a 3-manifold is called non-degenerate if the spectrum of the linearized Poincaré map associated to any (prime) closed Reeb orbit does not contain roots of unity when restricted to the contact structure. According to [18],

λ is said to be dynamically convex if $c_1(\ker \lambda)$ vanishes and the Conley-Zehnder index of every contractible closed orbit of the associated Reeb flow is at least 3. See § 2.1 for a definition of the index in 3-dimensions.

In [16, 17] it is proved that a closed connected tight contact 3-manifold M is the tight 3-sphere if, and only if, the contact structure can be realized as the kernel of a dynamically convex non-degenerate contact form admitting an unknotted closed Reeb orbit P with self-linking number -1 and Conley-Zehnder index 3. In fact, they show that the given orbit bounds a disk-like global surface of section for the Reeb flow, but much more can be said: there is an open book decomposition of M with disk-like pages and binding P , such that every page is a global surface of section. In particular, M is homeomorphic to S^3 . The following result from [20] states that the restriction on the Conley-Zehnder index can be dropped.

Theorem 1.7. *Let λ be a non-degenerate dynamically convex tight contact form on a closed connected 3-manifold M . A closed Reeb orbit P is the binding of an open book decomposition with disk-like pages which are global surfaces of section for the Reeb flow if, and only if, it is unknotted and has self-linking number -1 . In particular, M is homeomorphic to the 3-sphere when an orbit P with these properties exists.*

The geodesic flow restricted to the unit sphere bundle of a Finsler metric F coincides with the Reeb flow of the contact form α_F , as explained before. Suppose F is such a metric on S^2 satisfying (2), and assume γ is a closed geodesic whose lift $\dot{\gamma}$ is unknotted and has self-linking number -1 in SS^2 . Theorem 1.7 together with the statement below due to Harris and Paternain [13] provides, in the bumpy case, a contradiction to the existence of γ since the unit sphere bundle is not homeomorphic to the 3-sphere. Thus the weak flat knot type k_8 can not be realized by a closed geodesic in view of Lemma 1.4. The general case is discussed in Section 3.

Theorem 1.8 (Harris and Paternain). *If a Finsler metric F with reversibility r on S^2 is δ -pinched, for some $\delta > (r/(r+1))^2$, then α_F is dynamically convex.*

The proof of Theorem 1.8 relies on Rademacher's estimate for the length of closed geodesic loops, see § 2.4 below. As it will be clear, the proof of Theorem 1.5 shows that Harris-Paternain's pinching condition is sharp in the following sense: given $r \geq 1$ and $0 < \delta < (r/(r+1))^2$ there exists a δ -pinched Finsler metric on S^2 with reversibility r , such that α_F is not dynamically convex.

Remark 1.9. As the reader may already have noticed, our argument will actually show a possibly stronger statement than that of Theorem 1.5. In fact, if we assume (2) then the transverse knot $\dot{\gamma}/F(\dot{\gamma})$ associated to a closed prime geodesic γ can not be unknotted and have self-linking number -1 .

1.2.2. Convex energy levels C^2 -close to S^{2n-1} . The low-high action dichotomy in Theorem 1.6 above is, of course, an immediate consequence of the non-trivial analysis from [3].

Consider an unperturbed flow and a periodic orbit P with prime period T for which the linearized transverse Poincaré map is the identity. Then, roughly speaking, a prime closed orbit near P of a perturbed flow either has period $\sim T$, or has a very large period. This is a particular instance of Proposition 4.1 below which was extracted from [3]. The low-high action dichotomy is obtained when we take as the unperturbed flow the Reeb flow on S^{2n-1} induced by the contact form

$\lambda_0 = \frac{1}{2} \sum qdp - pdq$, since all orbits are periodic with the same period, and the transverse linearized Poincaré map is always the identity.

In the case $n = 2$ we study the relation between orbits with low and high action. Analyzing specific global behavior of the “round” flow on S^3 we are able to conclude that a short orbit P_s of the perturbed Reeb flow bounds a disk transverse to the flow. A long orbit P_l either stays far from P_s , and thus links many times with it, or gets close to P_s and again links many times since the linearized flow along P_s rotates almost uniformly.

2. PRELIMINARIES

This section is devoted to reviewing the definitions and facts necessary for the proofs that follow.

2.1. The Conley-Zehnder index in 3 dimensions. The Conley-Zehnder index is an invariant of the linearized dynamics along closed Reeb orbits, which we now describe in the 3-dimensional case.

Whenever $I \subset \mathbb{R}$ is a closed interval of length strictly less than $1/2$ satisfying $\partial I \cap \mathbb{Z} = \emptyset$, consider the integer $\hat{\mu}(I)$ defined by $\hat{\mu}(I) = 2k$ if $k \in I$, or $\hat{\mu}(I) = 2k + 1$ if $I \subset (k, k + 1)$. It can be extended to the set of all closed intervals of length strictly less than $1/2$ by $\hat{\mu}(I) = \lim_{\epsilon \rightarrow 0^+} \hat{\mu}(I - \epsilon)$.

Let α be a contact form on the 3-manifold N , inducing the contact structure $\xi = \ker \alpha$. Then ξ becomes a symplectic vector bundle with the bilinear form $d\alpha|_\xi$. Suppose $x : \mathbb{R} \rightarrow N$ is a contractible periodic trajectory of the Reeb vector R (uniquely defined by $i_R \alpha = 1$ and $i_R d\alpha = 0$) of period $T > 0$, and let $f : \mathbb{D} \rightarrow N$ be a map satisfying $f(e^{i2\pi t}) = x_T(t) := x(Tt)$. We can find a symplectic trivialization $f^* \xi \simeq \mathbb{D} \times \mathbb{R}^2$, which restricts to a trivialization $\Psi : x_T^* \xi \rightarrow S^1 \times \mathbb{R}^2$. If ϕ_t is the Reeb flow then $d\phi_t$ preserves ξ , and we get a smooth path of symplectic 2×2 matrices $t \in \mathbb{R} \mapsto \varphi(t) = \Psi_t \cdot d\phi_{Tt} \cdot \Psi_0^{-1}$, where Ψ_t is restriction of Ψ to $x_T^* \xi|_t$.

Given $w \in \mathbb{R}^2$, $w \neq 0$, define $\Delta(w) := \frac{1}{2\pi}(\vartheta(1) - \vartheta(0))$, where $\vartheta(t)$ is a continuous lift of the argument of $\varphi(t)w$. Then consider the closed real interval $I := \{\Delta(w) : w \in \mathbb{R}^2, w \neq 0\}$. It is easy to check that I has length strictly less than $1/2$. Following [19], one can define

$$(3) \quad \mu_{CZ}(x, T) = \hat{\mu}(I).$$

Note that $\mu_{CZ}(x, T) \geq 3$ if, and only if, $I \subset \{y \in \mathbb{R} : y > 1\}$. Once f is fixed, the above integer does not depend on the choice of symplectic trivialization $f^* \xi \simeq \mathbb{D} \times \mathbb{R}^2$. Nevertheless, the notation should indicate the dependence on the disk-map f , but $\mu_{CZ}(x, T)$ does not depend on f when $c_1(\xi)$ vanishes, see [19] for more details.

2.2. The self-linking number. Let (M, ξ) be a contact 3-manifold, $L \subset M$ be a knot transverse to ξ , and let $\Sigma \hookrightarrow M$ be a Seifert surface for L , that is, S is an orientable embedded connected compact surface $\Sigma \hookrightarrow M$ such that $L = \partial \Sigma$. Assume $\xi = \ker \lambda$ for some contact form λ . Since the bundle $\xi|_\Sigma$ carries the symplectic bilinear form $d\lambda$, there exists a smooth non-vanishing section Z of $\xi|_\Sigma$ which can be used to slightly perturb L to another transverse knot $L_\epsilon = \{\exp_x(\epsilon Z_x) : x \in L\}$. Here $\exp$ is any exponential map. A choice of orientation for Σ induces orientations of L and of L_ϵ . The *self-linking number* is defined as the

oriented intersection number

$$(4) \quad sl(L, \Sigma) := L_\epsilon \cdot \Sigma \in \mathbb{Z},$$

where M is oriented by $\lambda \wedge d\lambda$. It is independent of Σ when $c_1(\xi) \in H^2(M)$ vanishes.

2.3. Basics in Finsler geometry. We recall the basic definitions in Finsler geometry following [11]. See also [25, 8, 28]. The knowledgeable reader is encouraged to skip to § 2.4, and refer back only for the notation established here.

2.3.1. Connections and curvatures. Let $\pi : TM \rightarrow M$ be the tangent bundle of a manifold M , and denote $TM_0 := TM \setminus \{\text{zero section}\}$. Let VTM be the vertical subbundle $\ker d\pi \subset TTM$, with fiber $V_v TM$ over $v \in TM$. VTM_0 denotes its restriction to TM_0 . Whenever $(x^1, \dots, x^n)$ are coordinates on M we have natural coordinates

$$(x^1, \dots, x^n, y^1, \dots, y^n) \simeq \sum_i y^i \partial_{x^i}$$

on TM . Thus, $\{\partial_{y^1}, \dots, \partial_{y^n}\}$ is a local frame on VTM . On TM we have a vector field defined in natural coordinates by $C = \sum_{i=1}^n y^i \partial_{y^i}$, and the almost tangent structure $\mathcal{J}$, which is the VTM -valued 1-form on TM defined locally by $\mathcal{J} = \sum_{i=1}^n dx^i \otimes \partial_{y^i}$. There is a canonical linear isomorphism $i_v : T_{\pi(v)}M \simeq V_v TM$, for any given $v \in TM$, defined by $i_v(w) = \frac{d}{dt}|_{t=0} v + tw$. In natural coordinates: $i_v(w) = \sum_i w^i \partial_{y^i}$ if $w = \sum_i w^i \partial_{x^i}$. Thus $C_v = i_v(v)$.

A TTM -valued 1-form Γ on TM_0 satisfying

$$(5) \quad \Gamma^2 = I \quad \text{and} \quad \ker(\Gamma + I) = VTM_0$$

is a Grifone connection on M . In natural coordinates the equations $\Gamma \cdot \partial_{x^i} = \partial_{x^i} - 2\Gamma_i^j \partial_{y^j}$, $\Gamma \cdot \partial_{y^i} = -\partial_{y^i}$ define the connection coefficients Γ_i^j (we use Einstein summation convention). Considering the associated horizontal subbundle $HTM := \ker(\Gamma - I)$ we have a splitting

$$(6) \quad TTM = VTM \oplus HTM$$

and induced projections $P_V : TTM \rightarrow VTM$, $P_H : TTM \rightarrow HTM$. The isomorphisms $i_v^{-1} : V_v TM \simeq T_{\pi(v)}M$ and $d\pi : H_v TM \simeq T_{\pi(v)}M$ provide an isomorphism

$$(7) \quad (i_v^{-1} \circ P_V, d\pi \circ P_H) : T_v TM \simeq T_{\pi(v)}M \oplus T_{\pi(v)}M$$

when $v \neq 0$. If $\zeta = \delta x^i \partial_{x^i} + \delta y^i \partial_{y^i} \in T_v TM$ then

$$\zeta \simeq ((\delta y^i + \Gamma_k^i \delta x^k) \partial_{x^i}, \delta x^i \partial_{x^i}) \in T_{\pi(v)}M \oplus T_{\pi(v)}M$$

by the map (7).

The curvature form of Γ is the VTM -valued 2-form on TM_0 defined by

$$(8) \quad R(X, Y) = P_V([P_H(X), P_H(Y)])$$

where X, Y are vector fields on TM_0 .

Later we will need to consider lifts of a Grifone connection Γ . These are linear connections ∇ on VTM satisfying

$$(9) \quad HTM = \ker(X \mapsto \nabla_X C).$$

∇ is said to be symmetric if $\nabla_X \mathcal{J}(Y) - \nabla_Y \mathcal{J}(X) = \mathcal{J}([X, Y])$ for arbitrary vector fields X, Y on TM_0 . If ∇ has coefficients $\nabla_{\partial_{x^i}} \partial_{y^j} = \Gamma_{ij}^k \partial_{y^k}$ and $\nabla_{\partial_{y^i}} \partial_{y^j} = D_{ij}^k \partial_{y^k}$, in local natural coordinates, then this symmetry condition implies $D_{ij}^k = 0$, $\Gamma_{ij}^k = \Gamma_{ji}^k$ and $y^j \Gamma_{ij}^k = \Gamma_i^k$.

The curvature tensor of ∇ is

$$(10) \quad \tilde{R}(X, Y)Z = \nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X, Y]} Z$$

where X, Y are vector fields on TM_0 and Z is a section of VTM_0 . The curvature endomorphism of ∇ in the direction of $v \in TM_0$ is the linear map $R^v : T_{\pi(v)}M \rightarrow T_{\pi(v)}M$ defined by

$$(11) \quad R^v(u) = i_v^{-1}(\tilde{R}(v_h, u_h)i_v(v))$$

where $u_h, v_h \in T_v TM$ are the (unique) horizontal lifts of u, v , respectively.

2.3.2. Sprays and their geodesics. Recall that a spray is a continuous vector field S on TM , smooth on TM_0 , satisfying equations

$$(12) \quad i_S \mathcal{J} = C, \quad L_C S = S.$$

In local natural coordinates one can write $S = y^i \partial_{x^i} - 2G^i(x, y) \partial_{y^i}$. The G^i will be referred to as the spray coefficients, and they satisfy $G^i(x, ty) = t^2 G^i(x, y)$.

Every spray S defines a Grifone connection by $\Gamma_S := -L_S \mathcal{J}$. It follows from (12) that $L_C \Gamma_S = 0$ and that Γ_S is symmetric: if Γ_i^j are the connection coefficients in natural coordinates then $\partial_{y^k} \Gamma_i^j = \partial_{y^i} \Gamma_k^j$ for every i, j, k . Moreover, $\Gamma_i^j = \partial_{y^i} G^j$.

The set of symmetric lifts of Γ_S is non-empty: the Berwald connection of S is given in natural coordinates by

$$\nabla_{\partial_{y^i}} \partial_{y^j} = 0, \quad \nabla_{\partial_{x^i}} \partial_{y^j} = \Gamma_{ij}^k \partial_{y^k} \quad \text{where} \quad \Gamma_{ij}^k = \partial_{y^j} \Gamma_i^k = \frac{\partial^2 G^k}{\partial y^j \partial y^i}.$$

If $\zeta(t) \subset TM_0$ is an integral curve of S then $\zeta = \dot{\gamma}$ where $\gamma = \pi \circ \zeta$. This follows from (12). A curve $\gamma(t)$ on M is called a geodesic if $\dot{\gamma}(t)$ is an integral curve of S , where the condition $\dot{\gamma}(0) \neq 0$ is implicit.

Using Γ_S one defines the covariant derivative of a vector field $V(t)$ along a geodesic $\gamma(t)$. Namely, $\tilde{V}(t) := i_{\dot{\gamma}(t)}(V(t))$ defines a (vertical) vector field along the integral curve $\dot{\gamma}$ of S and, hence, the Lie derivative $L_S \tilde{V}$ is well-defined. We set

$$(13) \quad \frac{D_\gamma V}{dt} := i_{\dot{\gamma}(t)}^{-1}(P_V(L_S \tilde{V})).$$

In natural coordinates, if $V = V^i \partial_{x^i}$ and $\gamma^i(t) = x^i \circ \gamma$ then

$$(14) \quad \frac{D_\gamma V}{dt} = \left(\dot{V}^i + \Gamma_k^i V^k \right) \partial_{x^i}$$

where the Γ_k^i are evaluated at $(\gamma^1, \dots, \gamma^n, \dot{\gamma}^1, \dots, \dot{\gamma}^n)$. Note the drastic difference to the Riemannian case, where the Γ_k^i do not depend on $\dot{\gamma}^1, \dots, \dot{\gamma}^n$. Thus, in the more general present situation, covariant differentiation along arbitrary curves may not be defined only in terms of the spray, and other choices must be made.

Parallel transport $P_t : T_{\gamma(0)}M \rightarrow T_{\gamma(t)}M$ along a geodesic γ is defined by $P_t(V_0) := V(t)$, where $V_0 \in T_{\gamma(0)}M$, $\frac{D_\gamma V}{dt} = 0$ and $V(0) = V_0$.

Lemma 2.1. *Let ∇ be a symmetric lift of Γ_S . Then*

$$-R^v(u) = i_v^{-1}(R(S, u_h)) \quad \text{for any } u \in T_{\pi(v)}M$$

where $u_h \in H_v TM$ is the horizontal lift of u ($d\pi \cdot u_h = u$) and R is the curvature form of Γ_S . In particular, R^v is independent of the choice of the symmetric lift.

For completeness, and convenience of the reader, we include a proof of the above well-known standard fact in the appendix.

2.3.3. *The case of Finsler manifolds.* A Finsler metric on a manifold M is a continuous function $F : TM \rightarrow [0, +\infty)$, smooth on $TM_0 := TM \setminus \{\text{zero section}\}$ satisfying:

- (i) $F(tv) = tF(v), \forall v \in TM, \forall t > 0$. F is said to be positively homogeneous of degree 1.
- (ii) For each $v \in TM_0$ the quadratic form $g_v : T_{\pi(v)}M \times T_{\pi(v)}M \rightarrow \mathbb{R}$ given by

$$(15) \quad g_v(w_1, w_2) = \frac{1}{2} \frac{\partial^2}{\partial s \partial t} \Big|_{s=t=0} F^2(v + sw_1 + tw_2).$$

is positive definite. This is called the convexity condition.

The Legendre transform $\mathcal{L} : T^*M \rightarrow TM$ is the fiber-preserving homeomorphism defined in the following manner: given $\lambda \in T_p^*M$ then

$$(16) \quad \mathcal{L}(\lambda) := v^* \in T_pM \quad \text{where} \quad \lambda \cdot v^* - \frac{1}{2}F^2(v^*) = \sup_{v \in T_pM} \lambda \cdot v - \frac{1}{2}F^2(v).$$

Thus $\mathcal{L}(t\lambda) = t\mathcal{L}(\lambda)$, $\forall t \geq 0$ and $\lambda \in T^*M$, and $\lambda = 0 \Leftrightarrow \mathcal{L}(\lambda) = 0$. As a consequence, $\mathcal{L}$ is a diffeomorphism between $T^*M_0 := T^*M \setminus \{\text{zero section}\}$ and TM_0 . One defines the cometric

$$(17) \quad F^* : T^*M \rightarrow [0, +\infty), \quad F^* = F \circ \mathcal{L}.$$

Thus F^* is smooth on T^*M_0 , $F^*(t\lambda) = tF^*(\lambda) \forall t \geq 0$, and $F^*(\lambda) = 0 \Leftrightarrow \lambda = 0$.

On T^*M consider the tautological 1-form α_{taut} and the canonical symplectic structure $\Omega := d\alpha_{taut}$.

In natural coordinates $(x^1, \dots, p^1, \dots)$ on T^*M associated to a set of coordinates $(x^1, \dots, x^n)$ on M , $\alpha_{taut} = \sum_i p^i dx^i$ and $\Omega = \sum_i dp^i \wedge dx^i$. The Hamiltonian $H := \frac{1}{2}(F^*)^2$ induces the Hamiltonian vector field X_H by $-dH = \Omega(X_H, \cdot)$, and one checks that

$$(18) \quad S := \mathcal{L}_* X_H$$

is a spray. Its flow is the geodesic flow of the Finsler metric F , and S is called the geodesic spray.

If $g_{ij}(x, y)$ represents the quadratic form g_v (15) in natural coordinates, where $v \simeq (x, y)$, and if g^{ij} is the inverse of g_{ij} , then the geodesic spray coefficients are

$$(19) \quad G^i = \frac{1}{2} \gamma_{jk}^i y^j y^k \quad \text{where} \quad \gamma_{jk}^i := \frac{1}{2} g^{il} (\partial_{x^k} g_{lj} - \partial_{x^l} g_{jk} + \partial_{x^j} g_{kl}).$$

This is proved by analyzing the Euler-Lagrange equations of the variational problem associated to the integral $\int \frac{1}{2} F^2 dt$. The fundamental difference with Riemannian geometry is that all functions g^{il} , $\partial_{x^k} g_{lj}$, etc also depend on the fiber coordinates $y^1, \dots, y^n$, and not only on the $x^1, \dots, x^n$.

In the context of Finsler metrics, Γ_S has a symmetric lift which is more suitable than the Berwald connection. Consider the Cartan tensor: the $(0, 3)$ -tensor on the bundle VTM_0 defined in natural coordinates by

$$(20) \quad A_{ijk} = \frac{1}{4} \frac{\partial^3 F^2}{\partial y^i \partial y^j \partial y^k}.$$

Roughly speaking, it governs how g_v varies fiberwise. As explained in [25] or in [8], the Chern connection is the symmetric lift of Γ_S with coefficients

$$(21) \quad \Gamma_{ij}^k = \gamma_{ij}^k - g^{is} (A_{sjt} \Gamma_k^t - A_{jkt} \Gamma_s^t + A_{kst} \Gamma_j^t).$$

In the context of Finsler manifolds parallel transport has, as expected, useful metric properties.

Lemma 2.2. *Let $\gamma(t) \subset M$ be a geodesic, and let V, W be vector fields along γ . Then*

$$(22) \quad \frac{d}{dt}g_{\dot{\gamma}}(V, W) = g_{\dot{\gamma}}\left(\frac{D_{\dot{\gamma}}V}{dt}, W\right) + g_{\dot{\gamma}}\left(V, \frac{D_{\dot{\gamma}}W}{dt}\right).$$

In contrast to the Riemannian case, this formula may not hold when γ is not a geodesic. See the appendix for a proof.

A flag pole is a pair (Π, v) , where $v \in TM_0$ and Π is a 2-plane in $T_{\pi(v)}M$ containing v . In the context of Finsler manifolds explained above, the associated flag curvature is

$$(23) \quad K(\Pi, v) = \frac{g_v(-R^v(w), w)}{g_v(v, v)g_v(w, w) - g_v(v, w)^2}$$

where $w \in \Pi$ is any vector such that $\{v, w\}$ is linearly independent. In the Riemannian case, where all the tensors involved only depend on the base point, $K(\Pi, v)$ does not depend on $v \in \Pi$, and is the sectional curvature of Π .

A vector field J along a geodesic γ satisfying $\frac{D_{\dot{\gamma}}^2 J}{dt^2} - R^{\dot{\gamma}}(J) = 0$ is called a Jacobi field. This ODE is referred to as the Jacobi equation. The linearization of the geodesic flow can be suitably represented according to the following standard lemma. A proof is found in the appendix.

Lemma 2.3. *Let Φ_t be the flow of S and set $\zeta(t) = d\Phi_t \cdot \zeta(0)$ where $\zeta(0) \in T_{\dot{\gamma}(0)}TM$ is fixed. If J is the Jacobi field determined by $\zeta(0) \simeq (\frac{D_{\dot{\gamma}}J}{dt}(0), J(0))$ under the isomorphism (7) then*

$$(24) \quad \zeta(t) \simeq \left(\frac{D_{\dot{\gamma}}J}{dt}(t), J(t)\right).$$

The unit sphere bundle $SM = \{v \in TM : F(v) = 1\}$ has contact-type as a hypersurface inside TM_0 equipped with the symplectic structure $\Omega_F := (\mathcal{L}^{-1})^*\Omega$. This is so since C is a Liouville vector field, that is, $L_C\Omega_F = \Omega_F$. Thus $\alpha_F := i_C\Omega_F$ restricts to a contact form on SM . The geodesic spray S coincides with the Reeb vector field associated to $\alpha_F|_{SM}$, that is, it satisfies $i_S\alpha_F \equiv 1$, $i_S\Omega_F \equiv 0$. SM becomes a contact manifold with contact structure $\xi_F = \ker \alpha_F|_{SM} \subset TSM$.

Setting $(\mathbb{R}v)^\perp = \{w \in T_{\pi(v)}M \mid g_v(v, w) = 0\}$ then, since $g_v = \frac{1}{2} \frac{\partial^2 F^2}{\partial y^i \partial y^j} dx^i \otimes dx^j$ and $\alpha_F(v) = \frac{1}{2} \frac{\partial F^2}{\partial y^j} dx^j = \frac{1}{2} \frac{\partial^2 F^2}{\partial y^i \partial y^j} y^j dx^i = g_v(v, d\pi \cdot)$ in natural coordinates, one has

$$(25) \quad \xi_F|_v \simeq (\mathbb{R}v)^\perp \oplus (\mathbb{R}v)^\perp$$

under the isomorphism (7). Since the linearized flow $d\Phi_t$ preserves ξ_F we get, in view of (25) and (24), a familiar fact from Riemannian geometry: if $J(0), \frac{D_{\dot{\gamma}}J}{dt}(0) \in (\mathbb{R}\dot{\gamma}(0))^\perp$ then $J(t), \frac{D_{\dot{\gamma}}J}{dt}(t) \in (\mathbb{R}\dot{\gamma}(t))^\perp$, for every t .

2.4. Estimates on the Conley-Zehnder index. For the sake of completeness we quickly discuss the proof of Theorem 1.8. We use the notation established in the last paragraphs.

2.4.1. *A global symplectic trivialization of $\xi_F \subset SS^2$.* Let F be Finsler metric on S^2 , which is equipped with its orientation induced as a submanifold of $\mathbb{R}^3$. The unit sphere bundle SS^2 is equipped with the contact form α_F discussed in § 2.3.3. For every $v \in SS^2$, g_v is an inner-product on $T_{\pi(v)}S^2$, and there exists a unique vector $v^\perp \in T_{\pi(v)}S^2$ such that $\{v, v^\perp\}$ is a positively oriented g_v -orthonormal basis of $T_{\pi(v)}S^2$. Thus $(\mathbb{R}v)^\perp = \mathbb{R}v^\perp$ and we obtain a global trivialization

$$(26) \quad (v, (s, t)) \in SS^2 \times \mathbb{R}^2 \mapsto (sv^\perp, tv^\perp) \in (\mathbb{R}v)^\perp \oplus (\mathbb{R}v)^\perp \simeq \xi_F|_v$$

of ξ_F . The last identification is given by (25). One checks easily that this trivialization is symplectic with respect to $d\alpha_F$.

2.4.2. *Estimating the linearized twist.* Let $\gamma(t)$ be a geodesic on S^2 with unit speed, and choose $\zeta(0) \in \xi_F|_{\dot{\gamma}(0)} \subset T_{\dot{\gamma}(0)}SM$. Let Φ_t be the Reeb flow of α_F on SS^2 . Then, setting $\zeta(t) := d\Phi_t \cdot \zeta(0) \in \xi_F|_{\dot{\gamma}(t)}$, we can use (7) and identify $\zeta(t) \simeq (\frac{D_\gamma J}{dt}(t), J(t))$ as in (24), for some Jacobi field J . The vector field $\dot{\gamma}^\perp$ is parallel, as one can prove by using (22) and noting that $g_\gamma(\dot{\gamma}, \dot{\gamma}^\perp)$ and $g_\gamma(\dot{\gamma}^\perp, \dot{\gamma}^\perp)$ are constant in t . Writing $J = f\dot{\gamma}^\perp$ we get $\frac{D_\gamma J}{dt} = f'\dot{\gamma}^\perp$, $\frac{D_\gamma^2 J}{dt^2} = f''\dot{\gamma}^\perp$ and, consequently,

$$(27) \quad f'' = g_\gamma \left(\frac{D_\gamma^2 J}{dt^2}, \dot{\gamma}^\perp \right) = g_\gamma (R^\gamma(\dot{\gamma}^\perp), \dot{\gamma}^\perp) f = -K(T_\gamma S^2, \dot{\gamma}) f.$$

Abbreviating $K(T_\gamma S^2, \dot{\gamma})$ by $K(t)$ we get, after further identifying

$$\zeta(t) \simeq (\frac{D_\gamma J}{dt}(t), J(t)) \simeq (f', f) \doteq u(t)$$

via (26), an equation

$$\dot{u} = \begin{pmatrix} 0 & -K(t) \\ 1 & 0 \end{pmatrix} u.$$

Thus, if $\vartheta(t)$ is a smooth lift of the argument of $u(t)$ and the Finsler metric is positively curved then

$$(28) \quad \dot{\vartheta} \geq \delta \text{ for any } \delta \text{ satisfying } 0 < \delta < \min\{1, K_{\min}\},$$

where $K_{\min}$ is the infimum among all flag curvatures.

2.4.3. *Rademacher's estimate on the length of a closed geodesic.* Let l denote the infimum among all lengths of closed geodesic loops, and $K_{\max}$ be the supremum among all flag curvatures. The following estimate was obtained by Rademacher in [26], see [25] for a detailed account of the subject:

$$(29) \quad 0 < K_{\min} \leq K_{\max} \Rightarrow l \geq \frac{\pi(1 + \frac{1}{r})}{\sqrt{K_{\max}}}$$

where $r \geq 1$ is the reversibility. In the Riemannian case $r = 1$ and (29) is obtained from Klingenberg's estimate on the injectivity radius.

Lemma 2.4. *In the particular case of a positively curved Finsler metric on S^2 , every prime closed geodesic γ such that $\dot{\gamma}$ is contractible in SS^2 has length larger than or equal to $2\pi(1 + r^{-1})/\sqrt{K_{\max}}$.*

Proof. γ consists of at least two distinct closed loops since, otherwise, $\dot{\gamma}$ is not contractible in SS^2 . $\square$

2.4.4. *Estimating the index.* Consider a positively curved Finsler metric F with reversibility r that is strongly $(r/(r+1))^2$ -pinched. After rescaling we can assume

$$1 = K_{\min} \leq K_{\max} < \left(\frac{r+1}{r} \right)^2.$$

It follows by Lemma 2.4 and estimate (28) that ϑ varies strictly more than 2π along a closed geodesic γ with $\dot{\gamma}$ contractible in SS^2 , regardless of the choice of initial condition in $\ker \alpha_F|_{\dot{\gamma}(0)}$. According to (3) this proves Theorem 1.8.

3. PROOF OF THEOREM 1.5

We split the arguments in three parts. In § 3.1 we prove Lemma 1.4. In § 3.2 we use Theorem 1.7 to prove (i) in Theorem 1.5. Finally, in § 3.3 we exhibit for any $\epsilon > 0$ and $r \geq 1$, examples of Finsler metrics of Randers type on S^2 with reversibility r and flag curvatures in $\delta(r) - \epsilon < K \leq 1$ admitting geodesics with one transverse self-intersection.

3.1. Contact-topological invariants of k_8 . Let us denote by $\mathcal{B}_8 \subset \mathcal{B}$ the set of C^2 immersions of S^1 into S^2 with precisely one transverse self-intersection. Every $c \in \mathcal{B}_8$ induces an embedded copy of S^1 inside $F^{-1}(1) \subset TS^2$ given by $\dot{c}/F(\dot{c})$. Recall the Legendre transform $\mathcal{L}_F : T^*S^2 \rightarrow TS^2$ induced by F , which is a continuous fiber-preserving map that restricts to a smooth diffeomorphism $T^*S^2 \setminus 0 \simeq TS^2 \setminus 0$ and induces a cometric $F^* = F \circ \mathcal{L}_F$. Then the tautological 1-form α_{taut} restricts to a contact form on $(F^*)^{-1}(1)$. Consequently $\alpha_F := (\mathcal{L}_F^{-1})^* \alpha_{\text{taut}}$ restricts to a contact form on $F^{-1}(1)$, and clearly $\dot{c}/F(\dot{c})$ is positively transverse to the contact structure $\xi_F := \ker \alpha_F$. It is not hard to check that any two $c_0, c_1 \in \mathcal{B}_8$ are homotopic through curves $c_s \in \mathcal{B}_8$, $s \in [0, 1]$, so that we get a corresponding isotopy $\dot{c}_s/F(\dot{c}_s)$ through knots which are transverse to ξ_F , thus preserving the knot type and the self-linking number. Consequently, it suffices to exhibit one element $c \in \mathcal{B}_8$ such that $\dot{c}/F(\dot{c})$ is unknotted and has self-linking number -1 .

First we prove Lemma 1.4 for the metric $F_0(v) := \sqrt{g_0(v, v)}$, where g_0 is the Riemannian metric induced by isometrically embedding S^2 in $\mathbb{R}^3$ as the unit sphere. Taking polar coordinates (θ, R) in $\mathbb{C}$, $\theta \in \mathbb{R}/2\pi\mathbb{Z}$ and $R \geq 0$, consider the embedding

$$\gamma : \mathbb{C} \rightarrow S^3 \subset \mathbb{C}^2; (\theta, R) \mapsto \frac{1}{\sqrt{1+R^2}}(1, Re^{i\theta}).$$

Denote by

$$\lambda_0 = \frac{1}{4i}(\bar{z}dz - zd\bar{z} + \bar{w}dw - wd\bar{w}) = \frac{1}{2}(q_0dp_0 - p_0dq_0 + q_1dp_1 - p_1dq_1)$$

the Liouville form on $\mathbb{C}^2 \simeq \mathbb{R}^4$ with complex coordinates $(z = q_0 + ip_0, w = q_1 + ip_1)$. The embedded circles $\gamma_R = \gamma(\cdot, R)$ converge, as $R \rightarrow \infty$, to the Hopf fiber $P_0 = \{\theta \mapsto (0, e^{i\theta}) \mid \theta \in [0, 2\pi]\}$ in the C^1 -topology. All γ_R are positively transverse (with respect to λ_0) to $\ker \lambda_0$. It is well known that $\text{sl}(P_0) = -1$, which implies $\text{sl}(\gamma_R) = -1 \forall R > 0$. Moreover, γ_{R_0} is clearly unknotted since it is the boundary of the embedded disk $\mathcal{D}_{R_0}$ parametrized by $\gamma|_{\{R \leq R_0\}}$.

Identifying $S^3 \simeq SU(2)$ via

$$(z, w) \simeq \begin{pmatrix} z & w \\ -\bar{w} & \bar{z} \end{pmatrix}, \quad |z|^2 + |w|^2 = 1,$$

and considering the matrices

$$j = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \quad \text{and} \quad k = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}$$

there is a double cover $D : S^3 \rightarrow F_0^{-1}(1)$ given by

$$A \in SU(2) \simeq S^3 \mapsto (A^{-1}jA, -A^{-1}kA) \in F_0^{-1}(1)$$

where we see a unit vector $(x, y, t) \in S^2 \subset \mathbb{R}^3$ sitting inside S^3 as (it, u) where $u = x + iy$. Here $A^{-1}jA$ represents the base point and $-A^{-1}kA$ represents the tangent vector. We have $D^*\alpha_{F_0} = 4\lambda_0|_{S^3}$ (cf. [10, 13]). The factor 4 appears since a Hopf circle on S^3 , which has λ_0 -action equal to π , projects onto the unit velocity vector of a great circle prescribed twice, which has length 4π .

The group of deck transformations of D is precisely $\{id, a\}$, where a is the antipodal map. Thus, for each $R_0 > 0$, $\Gamma_{R_0} := D \circ \gamma_{R_0}$ is an embedded knot in $F_0^{-1}(1)$ since the curve γ_{R_0} does not contain pairs of antipodal points. It is clearly transverse to ξ_{F_0} . By the same token, $\tilde{\mathcal{D}}_{R_0} := D(\mathcal{D}_{R_0})$ is an embedded disk with boundary Γ_{R_0} , proving that Γ_{R_0} is unknotted. Moreover, since D is a 1-1 contactomorphism of a neighborhood of $\mathcal{D}_{R_0}$ in S^3 onto a neighborhood of $\tilde{\mathcal{D}}_{R_0}$ in $F_0^{-1}(1)$, we get that the self-linking number of Γ_{R_0} is also -1 . We concluded that each Γ_{R_0} is unknotted and has self-linking number -1 in the contact manifold $(F_0^{-1}(1), \xi_{F_0})$.

It only remains to find $c \in \mathcal{B}_8$ such that Γ_1 is transversely isotopic to $\dot{c}/F_0(\dot{c})$. Let $c : [0, 2\pi] \rightarrow S^2$ be defined by the equation

$$c(\theta) = \pi \circ \Gamma_1(\theta)$$

where $\pi : TS^2 \rightarrow S^2$ is the bundle projection. It is clear from the formula

$$c(\theta) = \frac{1}{2}(1 + \cos 2\theta, \sin 2\theta, 2 \sin \theta) \in \mathbb{R}^3$$

that $c \in \mathcal{B}_8$. Since Γ_1 is positively transverse to ξ_{F_0} , we have

$$\alpha_{F_0}|_{\Gamma_1(\theta)} \cdot \dot{\Gamma}_1(\theta) = g_0(c(\theta))(\dot{c}(\theta), \Gamma_1(\theta)) > 0 \quad \forall \theta.$$

Thus we find a C^1 lift $\vartheta(\theta) \in (-\frac{\pi}{2}, \frac{\pi}{2})$ for the g_0 -angle between Γ_1 and $\dot{c}/F_0(\dot{c})$, and can define a transverse homotopy between Γ_1 and $\dot{c}/F_0(\dot{c})$ keeping the base points fixed by the formula

$$h_s(\theta) = \frac{(1-s)\frac{\dot{c}(\theta)}{F_0(\dot{c}(\theta))} + s\Gamma_1(\theta)}{F_0\left((1-s)\frac{\dot{c}(\theta)}{F_0(\dot{c}(\theta))} + s\Gamma_1(\theta)\right)} \in T_{c(\theta)}S^2 \cap F_0^{-1}(1).$$

It remains to show that $\{h_s\}_{s \in [0,1]}$ is a transverse isotopy. The only possibility for self-intersections of the curves $\theta \mapsto h_s(\theta) \in F_0^{-1}(1)$ is at the values $\theta = 0$ and $\theta = \pi$ where the curve $c(\theta)$ self-intersects at the point $(1, 0, 0)$. Looking at the formulas

$$\begin{aligned} \dot{c}(\theta) &= (-\sin 2\theta, \cos 2\theta, \cos \theta) \in T_{c(\theta)}S^2 \\ \Gamma_1(\theta) &= (-\tfrac{1}{2}\sin 2\theta, \tfrac{1}{2}(\cos 2\theta - 1), \cos \theta) \in T_{c(\theta)}S^2 \end{aligned}$$

we note that both $\dot{c}(0)$ and $\Gamma_1(0)$ point at north hemisphere, while $\dot{c}(\pi)$ and $\Gamma_1(\pi)$ point at the south hemisphere. Thus the formula for h_s does not produce self-intersections and, consequently, is a transverse isotopy. Lemma 1.4 is proved for the metric F_0 .

Now we consider a general Finsler metric $F : TS^2 \rightarrow [0, +\infty)$. We have associated Legendre transforms $\mathcal{L}_{F_0}, \mathcal{L}_F$ and cometrics $F_0^* = F_0 \circ \mathcal{L}_{F_0}$, $F^* =$

$F \circ \mathcal{L}_F$. The map $\Psi : (F^*)^{-1}(1) \rightarrow (F_0^*)^{-1}(1)$ defined by $\Psi(p) = p/F_0^*(p)$ satisfies $\Psi^*(\alpha_{\text{taut}}|_{(F_0^*)^{-1}(1)}) = \frac{1}{F_0^*} \alpha_{\text{taut}}|_{(F^*)^{-1}(1)}$, so that it is a contactomorphism. Hence we get a contactomorphism

$$\Phi := \mathcal{L}_{F_0} \circ \Psi \circ \mathcal{L}_F^{-1} : F^{-1}(1) \rightarrow F_0^{-1}(1).$$

Given any immersion $c : S^1 \rightarrow S^2$ we construct two transverse embeddings $V_c, W_c : S^1 \rightarrow F_0^{-1}(1)$ covering c as follows: $V_c = \dot{c}/F_0(\dot{c})$ and $W_c = \Phi(\dot{c}/F(\dot{c}))$. Since Φ preserves co-orientations induced by α_F and α_{F_0} , we have

$$g_0(c(\theta))(V_c(\theta), W_c(\theta)) > 0, \forall \theta \in \mathbb{R}/2\pi\mathbb{Z} \simeq S^1.$$

Now let $c_0 \in \mathcal{B}_8$ be arbitrary. Assume its self-intersection point is $c_0(0) = c_0(\pi)$. We clearly can find a homotopy $c_s : S^1 \rightarrow S^2$, $s \in [0, 1]$, starting at c_0 , such that $c_s \in \mathcal{B}_8$ for every $s \in [0, 1]$, and the immersion c_1 has a unique self-intersection point at $c_1(0) = c_1(\pi)$ satisfying $\dot{c}_1(0) = -\dot{c}_1(\pi)$ (negative self-tangency). This induces corresponding isotopies $V_{c_s}, W_{c_s} : S^1 \rightarrow F_0^{-1}(1)$ through transverse knots. Now define a homotopy $H_\tau : S^1 \rightarrow F_0^{-1}(1)$ satisfying $H_0 = V_{c_1}$ and $H_1 = W_{c_1}$ by

$$H_\tau(\theta) = \frac{(1-\tau)V_{c_1}(\theta) + \tau W_{c_1}(\theta)}{F_0((1-\tau)V_{c_1}(\theta) + \tau W_{c_1}(\theta))},$$

where $\tau \in [0, 1]$. The map H_τ is well-defined since $g_0(c_1(\theta))(V_{c_1}(\theta), W_{c_1}(\theta)) > 0$, for all θ , and hence the denominator above never vanishes. However, there could be some value of τ where H_τ is not a knot in $F_0^{-1}(1)$. This would only be the case if H_τ has self-intersections, which is only possible at the values $\theta = 0$ and $\theta = \pi$. Note, however, that this never happens because of the condition $\dot{c}_1(0) = -\dot{c}_1(\pi)$. We succeeded in showing that the knots V_{c_0} and W_{c_0} are transversally isotopic in $(F_0^{-1}(1), \xi_{F_0})$. We already showed before that V_{c_0} is unknotted and has self-linking number -1 . Thus the same is true for the knot $\dot{c}/F(\dot{c}) \subset (F^{-1}(1), \xi_F)$.

3.2. Non-existence of geodesics. Let F be a Finsler metric on S^2 satisfying (2). Then, as remarked in § 2.3.3, the pull-back of the tautological 1-form to TTS^2 , via the inverse Legendre transform induced by F , restricts to a contact form λ on SS^2 , which induces the contact structure $\xi := \ker \lambda$. By Theorem 1.8, λ is dynamically convex.

Assume, by contradiction, that there exists a prime closed geodesic γ with unit speed, such that $P = \{t \mapsto \dot{\gamma}(t)\} \subset SS^2$ is a closed unknotted Reeb orbit with self-linking number -1 .

In the case λ is non-degenerate Theorem 1.7 implies that SS^2 is homeomorphic to S^3 , a contradiction. It remains to consider the degenerate case. Denote by

$$\mathcal{O}_\lambda = \{f \in C^\infty(SS^2, \mathbb{R}^+) \mid f\lambda \text{ is non-degenerate}\}.$$

The following lemma, which we state without proof, is an adaptation of Lemma 6.8 in [18] to our situation, see also [15]. The proof is straightforward.

Lemma 3.1. *There exists a sequence $f_k \in \mathcal{O}_\lambda$ converging to 1 in the C^∞ -topology as $k \rightarrow +\infty$, such that each contact form $\lambda_k := f_k \lambda$ admits $P = (x, T)$ as a closed Reeb orbit.*

We will prove now that λ_k is dynamically convex for all k sufficiently large. We denote by ϕ_t the Reeb flow of λ and by ϕ_t^k the Reeb flow of λ_k .

Consider the global $d\lambda$ -symplectic trivialization $\Psi : \xi \xrightarrow{\sim} SS^2 \times \mathbb{C}$ described in (26). Since $\lambda_k \rightarrow \lambda$ in C^∞ we can find $d\lambda_k$ -symplectic trivializations $\Psi_k : \xi \xrightarrow{\sim}$

$SS^2 \times \mathbb{C}$ such that $\Psi_k \rightarrow \Psi$ in C^∞ . Given $p_0 \in SS^2$ and $v(0) \in \xi_{p_0} \setminus 0$ arbitrary, the solution $v(t) = d\phi_t \cdot v(0) \in \xi_{\phi_t(p_0)}$ of the linearized λ -Reeb flow can be represented using the frame Ψ as a curve $v(t) \simeq r(t)e^{i\theta(t)} \subset \mathbb{C}$, where $r(t) > 0$ and $\theta(t) \in \mathbb{R}$ is any continuous lift of the argument. There exists $a > 0$ such that

$$(30) \quad \dot{\theta}(t) > a,$$

for all t , independently of the choice of p_0 and $v(0)$. The existence of a follows from estimate (28). Moreover, if p_0 is a point on a closed contractible λ -Reeb orbit $P = (\bar{x}, \bar{T})$ then

$$(31) \quad \theta(\bar{T}) - \theta(0) > 2\pi.$$

This is a consequence of (3) and of the dynamical convexity of λ . In the following, solutions of the linearized λ_k -Reeb flow will be represented similarly by curves in the complex plane with the use of the frame Ψ_k .

Arguing indirectly, suppose there exists a subsequence of f_k , again denoted by f_k , such that λ_k is not dynamically convex. Then there exists a contractible closed λ_k -Reeb orbit $P_k = (x_k, T_k)$, with $\mu_{CZ}(P_k) \leq 2$, for each k . Assume first that $T_k \rightarrow +\infty$ as $k \rightarrow +\infty$. Since $\lambda_k \rightarrow \lambda$ and $\Psi_k \rightarrow \Psi$ in the C^∞ -topology, and SS^2 is compact, inequality (30) holds for any linearized solution $v_k \simeq r_k e^{i\theta_k}$ of the λ_k -Reeb flow over P_k , if k is large. In view of the geometric definition of the Conley-Zehnder index explained in § 2.1, this implies

$$\mu_{CZ}(P_k) \geq \frac{\theta_k(T_k) - \theta_k(0)}{\pi} - 1 > \frac{aT_k}{\pi} - 1 \rightarrow +\infty$$

as $k \rightarrow \infty$, in contradiction with $\mu_{CZ}(P_k) \leq 2$.

Now assume that $T_k = \int_{P_k} \lambda_k$ has a bounded subsequence. By the Arzelà-Ascoli theorem we find a converging subsequence, still denoted T_k , such that $T_k \rightarrow T_0$, $x_k(T_k \cdot) \rightarrow x_0(T_0 \cdot)$ in C^∞ where $P_0 = (x_0, T_0)$ is a closed λ -Reeb orbit. We also find a solution $v_k = r_k e^{i\theta_k}$ of the linearized flow over P_k , with $v_k(0)$ bounded and bounded away from $0 \in \mathbb{C}$, satisfying $\theta_k(T_k) - \theta_k(0) \leq 2\pi$ for each k . Again using that $\lambda_k \rightarrow \lambda$ and $\Psi_k \rightarrow \Psi$ in C^∞ , a subsequence of v_k converges (in C^∞) to a solution $v_0 = r_0 e^{i\theta_0}$ over P_0 satisfying $\theta_0(T_0) - \theta_0(0) \leq 2\pi$, in contradiction to (31).

Therefore we end up with a sequence λ_k converging to λ in the C^∞ -topology such that, for each k large enough, λ_k is a dynamically convex non-degenerate tight contact form on SS^2 admitting the unknotted closed orbit P with $sl(P) = -1$. Reasoning as before, Theorem 1.7 leads to the contradiction $SS^2 \simeq S^3$. The proof of (i) in Theorem 1.5 is complete.

3.3. Examples of Randers type. Here we prove (ii) in Theorem 1.5. A Riemannian metric a and a 1-form b on a manifold M induce a Randers metric

$$(32) \quad F(v) = \sqrt{a(v, v)} + b(v), \quad v \in TM,$$

precisely when $|b|_a < 1$ everywhere. These form an interesting and rich family of Finsler geometries, vastly studied in the literature.

3.3.1. Zermelo navigation. A pair (h, X) , where h is a Riemannian metric on M and X is a vector field satisfying $\sqrt{h(X, X)} < 1$, is called a Zermelo navigation data. It induces a Randers-type metric on T^*M by $F^*(\lambda) = \sqrt{h^*(\lambda, \lambda)} + \lambda \cdot X$, where h^* is the dual of h . The pull-back F of F^* by the Legendre transform is a Finsler metric on M , which is said to solve the associated Zermelo navigation

problem. In fact, its geodesic flow parametrizes the movement of a particle on M under the additional influence of a tangential wind, see [8] for a detailed discussion.

Remark 3.2. It is curious that F is of Randers type, that is, Legendre transformation preserves the form of the metric. To see this, consider $(x^1 \dots x^n, y^1 \dots y^n)$ natural coordinates on TM , with dual coordinates $(x^1 \dots x^n, p_1 \dots p_n)$ on T^*M . Then $F^* = \sqrt{h^{rs}p_r p_s} + X^k p_k$ and

$$y^i = \frac{1}{2} \frac{\partial (F^*)^2}{\partial p_i} = \left(\frac{h^{ij} p_j}{\sqrt{h^{rs} p_r p_s}} + X^i \right) F^* \Rightarrow p_k = h_{ki} (y^i - F X^i) \frac{\sqrt{h^{rs} p_r p_s}}{F}.$$

Plugging into the formula for $F = F^*$, and writing $\Delta = p_k X^k$, we get

$$F = \frac{F - \Delta}{F} \sqrt{h_{kl} (y^k - F X^k) (y^l - F X^l)} + \Delta$$

which gives $F = \sqrt{h_{kl} (y^k - F X^k) (y^l - F X^l)}$. Raising to the square and expanding the right side we get a second degree polynomial $\epsilon F^2 + BF + C = 0$, with $\epsilon = (1 - X_l X^l)$, $B = 2y^k X_k$ and $C = -h_{kl} y^k y^l$. Here we lowered the indices of X with the metric h . Solving we get $F = \sqrt{a_{ij} y^i y^j} + b_k y^k$, with $a_{ij} = \epsilon^{-1} h_{ij} + \epsilon^{-2} X_i X_j$ and $b_k = -\epsilon^{-1} X_k$.

The behavior of the geodesic flow of F is better understood if we work on T^*M equipped with its canonical symplectic structure $\Omega \simeq dp_i \wedge dx^i$ and Hamiltonian $\frac{1}{2}(F^*)^2$. This discussion is based on [25]. We can write

$$F^* = H + K \quad \text{with} \quad H = \sqrt{h^{ij} p_i p_j} \quad K = p_i X^i.$$

The Hamiltonian vector fields are $X_{F^*} = X_H + X_K$, where $i_{X_H} \Omega = -dH$ and $i_{X_K} \Omega = -dK$. If R_t is the flow of X then the flow of X_K is $\lambda \mapsto (dR_t^{-1})^* \cdot \lambda$. If X is Killing with respect to the metric h then $\{H, K\} = 0$ since R_t are isometries. Thus the flows Ψ_t^H and Ψ_t^K of X_H and X_K respectively, commute and, consequently, $\Psi_t^{F^*} = \Psi_t^H \circ \Psi_t^K$. The geodesic flow is precisely the Hamiltonian flow $\Psi_t^{(F^*)^2/2}$ of $\frac{1}{2}(F^*)^2$. Hence

$$(33) \quad \Psi_t^{(F^*)^2/2} = \Psi_t^{F^*} = \Psi_t^H \circ \Psi_t^K = \Psi_t^K \circ \Psi_t^H \quad \text{on the unit sphere bundle}$$

since $X_{(F^*)^2/2} = F^* X_{F^*}$. Since H is constant along trajectories of $X_{H^2/2}$ we have $\Psi_t^H(\lambda) = \Psi_{t/H(\lambda)}^{H^2/2}(\lambda)$, which we can use to finally arrive at

$$(34) \quad \Psi_t^{(F^*)^2/2}(\lambda) = \Psi_t^K \circ \Psi_{t/H(\lambda)}^{H^2/2}(\lambda), \quad \forall \lambda \in (F^*)^{-1}(1).$$

Consider the Legendre transform $\mathcal{L}(\lambda) \in T_x M$ associated to $\frac{1}{2}(F^*)^2$, where $\lambda = p_i dx^i \in T_x^* M$ is some covector. If $F^*(\lambda) = 1$ we have

$$\mathcal{L}(\lambda) = \sum_i \frac{1}{2} (\partial_{p_i} (F^*)^2) \partial_{x^i} = \left(\frac{h^{ij} p_j}{\sqrt{h^{rs} p_r p_s}} + X^i \right) \partial_{x^i} = \mathcal{L}_h \left(\frac{\lambda}{H(\lambda)} \right) + X$$

where $\mathcal{L}_h$ is the Legendre transform associated to $\frac{1}{2}H^2$. Thus, setting $\lambda = \mathcal{L}^{-1}(v)$, we find

$$(35) \quad \mathcal{L}_h \left(\frac{\lambda}{H(\lambda)} \right) = v - X, \quad \forall v \in SM = F^{-1}(1).$$

Fix $v \in SM$ and let $c(t)$ be the geodesic of F satisfying $\dot{c}(0) = v$. Then $c(t)$ is the base point of $\Psi_t^{(F^*)^2/2}(\lambda)$. The base point of $\Psi_{t/H(\lambda)}^{H^2/2}(\lambda)$ is equal to $\gamma_0(t)$, where γ_0

is the geodesic with respect to the metric h satisfying $\dot{\gamma}_0(0) = \mathcal{L}_h(\lambda/H(\lambda)) = v - X$, by (35). This fact and (34) imply

$$(36) \quad c(t) = R_t(\gamma_0(t)).$$

This formula will be used later. The reversibility of F is

$$(37) \quad r = \sup_{x \in M} \frac{1 + |X(x)|_h}{1 - |X(x)|_h}.$$

3.3.2. Pinched surfaces of revolution. Let (x, y, z) be standard Euclidean coordinates in 3-space. We consider a surface S of revolution around the z -axis symmetric under reflection with respect to the xy -plane, with the Riemannian metric induced by the ambient Euclidean metric. Denoting $\rho = \sqrt{x^2 + y^2}$, then S is determined by a curve in the ρz -plane, which we assume parametrized by a parameter s satisfying $\dot{\rho}^2 + \dot{z}^2 = 1$. We make $s = 0$ correspond to the equator, where the radius is $R = \rho(0)$. We always assume $R < 1$.

The symmetry condition forces ρ to be an even function of s in its domain $(-L, L)$. If S is a sphere then $L < +\infty$, $\dot{\rho} \rightarrow -1$ as $\rho \rightarrow \pm L$, the length of a meridian is $2L$ and S intersects the z -axis in two poles. In the complement of the poles we have obvious coordinates $(s, \theta) \in (-L, L) \times \mathbb{R}/2\pi\mathbb{Z}$.

The Gaussian curvature is $K = -\ddot{\rho}/\rho$, and we denote by K_{max} and K_{min} its maximum and minimum, respectively. If K is everywhere positive then the maximal radius is attained at the equator, and the maximal height is attained at the poles. We wish to construct S in a way that $K \equiv K_{min} = 1$ holds at the equator. Then $\ddot{\rho}(0) = -R$ and, moreover, $K_{max} \geq 1/R^2$ is a necessary condition. In fact, using $\dot{\rho}(0) = 0$, we compute

$$\begin{aligned} 1 - R^2 &= \int_0^L (\rho^2 + \dot{\rho}^2)' ds = \int_0^L (K - 1)(-2\rho\dot{\rho}) ds \\ &\leq (K_{max} - 1) \int_0^L (-2\rho\dot{\rho}) ds = (K_{max} - 1)R^2. \end{aligned}$$

Now we claim that if $K_{max} > 1/R^2$ then S with all the above properties exists. To see that, consider a smooth function $g : [0, R^2] \rightarrow [0, +\infty)$ satisfying $g(0) = 1$, $g(R^2) = 0$, $g' \in [-K_{max}, -1]$, $g' \equiv -K_{max}$ near 0, $g' \equiv -1$ near R^2 and $g'' \geq 0$. Note that $g(x) = R^2 - x$ for $x \sim R^2$. Let $\rho(s)$ be the unique solution of

$$(38) \quad \dot{\rho} = -\sqrt{g(\rho^2)}, \quad \rho(0) = R$$

for $s \geq 0$, which coincides² with $R \cos s$ for s small. Then $\dot{\rho}(0) = 0$ and there exists $L > 0$ such that $\rho(s) \rightarrow 0^+$ and $\dot{\rho}(s) \rightarrow -1^+$ as $s \rightarrow L^-$. This solution ρ determines a C^1 -embedded disk in the half-space $z \geq 0$, which can be reflected to provide the required C^1 sphere of revolution S . One can check that S is smooth, that the maximal value of the Gaussian curvature is K_{max} (attained around poles), and that the minimal value $K_{min} = 1$ (attained around the equator).

Note that $g(x) \geq R^2 - x \forall x \in [0, R^2]$. This implies

$$(39) \quad \rho(s) \leq R \cos s, \quad \forall s \in [0, L].$$

²Here it should be noted that (38) does not have a unique solution, as one can see by considering the constant function R .

3.3.3. Estimating a return time. Fix any $R \in (0, 1)$ and let $g : [0, R^2] \rightarrow [0, +\infty)$ be a function as in §§ 3.3.2 with $K_{max} > R^{-2}$. Consider the associated unique solution $\rho(s) : [0, L] \rightarrow \mathbb{R}$ of (38) which equals $R \cos s$ for small values of s . We claim that, for any $b > 1$, it is possible to make $2L < b\pi R$ by taking K_{max} close enough to R^{-2} .

To prove this, let $h : [0, R^2] \rightarrow [0, +\infty)$ be the continuous function defined by $h(x) = 1 - K_{max}x$ if $x \in [0, x_*]$ and $h(x) = R^2 - x$ if $x \in [x_*, R^2]$, where $x_* = (1 - R^2)/(K_{max} - 1) \in (0, R^2)$. It is imediate that $h(x) = g(x)$ for all x in a neighborhood of $\{0, R^2\}$. Since $g'' \geq 0$, we have $g \geq h$.

Let $\xi(s)$ be unique solution of $\dot{\xi} = -\sqrt{h(\xi^2)}$ with initial condition $\xi(0) = R$ coinciding with $R \cos s$ when $s > 0$ is small. We want to estimate the first $s_* > 0$ such that $\xi(s_*) = 0$. Observe that ξ satisfies $\ddot{\xi} = -K_{max}\xi$ if $0 \leq \xi \leq \sqrt{x_*}$ and satisfies $\ddot{\xi} = -\xi$ if $\sqrt{x_*} \leq \xi \leq R$. This implies that $s_* = s_1 + s_2$ where s_1 is such that $\xi(s_1) = \sqrt{x_*}$.

Now we prove that if we choose K_{max} sufficiently close to $1/R^2$ then s_* is close to $\frac{\pi R}{2}$. Observe that $\xi(s) = R \cos s$ for $0 \leq s \leq s_1$. Thus if $K_{max} \rightarrow R^{-2}$, $x_* \rightarrow R^2$ and, therefore, $s_1 \rightarrow 0$. For $s \geq s_1$, $\xi(s)$ is a solution of $\ddot{\xi} = -K_{max}\xi$ satisfying $\xi(s_1) = \sqrt{x_*}$ and $\dot{\xi}(s_1) < 0$. Thus the time s_2 that it takes to reach zero is smaller than the time $R \cos(\sqrt{K_{max}}s)$ takes to decay from R to 0, which is $\frac{\pi}{2\sqrt{K_{max}}}$. Consequently

$$s_2 < \frac{\pi}{2\sqrt{K_{max}}} \rightarrow \frac{\pi R}{2}$$

as $K_{max} \rightarrow 1/R^2$. Thus $K_{max} \sim R^{-2}$ implies $s_* = s_1 + s_2 < b\pi R/2$.

To estimate the length of the meridian observe that $\rho(s) \leq \xi(s)$ for all s since $g \geq h$ on $[0, R^2]$. Thus the length $2L$ of the meridian is at most $2s_*$ which is smaller than $b\pi R$ for K_{max} close enough to R^{-2} .

3.3.4. Introducing the wind and completing the proof. So far we have not fixed any of the data explicit in the statement of the Theorem 1.5.

Let $r \geq 1$ be given. Consider $\epsilon > 0$ small and numbers R, K_{max} satisfying

$$(40) \quad \frac{r}{r+1} - \epsilon < R < \frac{r}{r+1}, \quad \left(\frac{r}{r+1} \right)^{-2} < \frac{1}{R^2} < K_{max} < \left(\frac{r}{r+1} - \epsilon \right)^{-2}.$$

Following the construction in §§ 3.3.2, we can find a smooth surface of revolution (S, h) with Gaussian curvature taking values in $[1, K_{max}]$, and with an equator of radius R . We can arrange so that the curvature equals K_{max} near the poles, and equals 1 near the equator.

By the discussion of §§ 3.3.3 we can assume, after making ϵ small enough, that the length T of a meridian satisfies

$$(41) \quad T \leq \left(\frac{r+1}{r} \right) \pi R < \pi.$$

We can also assume that $(\frac{r-1}{r+1})/(\frac{r}{r+1} - \epsilon) \sim \frac{r-1}{r} < 1$ by making ϵ even smaller. Take $\eta \geq 0$ so that $\eta R = \frac{r-1}{r+1}$. Similarly to [21], consider the vector field

$$(42) \quad X = \eta \frac{\partial}{\partial \theta}$$

and let F be the Randers metric on S induced by the navigation data (h, X) , as explained in § 3.3.1. By (37) F has reversibility r . Later we shall need to note that

$$(43) \quad T\eta \leq \pi \left(\frac{r-1}{r+1} \right) \left(\frac{r}{r+1} - \epsilon \right)^{-1} < \pi$$

Crucial to our analysis is the fact that all flag curvatures of F are independent of the chosen flagpole and coincide with the Gaussian curvatures of h , see [8].

Lemma 3.3. *Let x be a point in the equator and let $v \in T_x S$ satisfy $F(v) = 1$ and $h(v - X, X) > 0$ (≥ 0). Then the geodesic $c(t)$ with respect to F with initial condition $\dot{c}(0) = v$ satisfies $h(\dot{c}(t), X \circ c(t)) > 0$ (≥ 0), $\forall t \geq 0$.*

Proof. Clearly X is Killing for h . According to (36), $c(t) = R_t \circ \gamma_0(t)$, where R_t is the flow of X and γ_0 is a geodesic with respect to h with $\dot{\gamma}_0(0) = v - X$. Thus, in view of the Clairaut integral for surfaces of revolution, $h(\dot{\gamma}_0, X \circ \gamma_0) > 0$ (≥ 0) for $t \geq 0$. We can estimate

$$\begin{aligned} h(\dot{c}, X \circ c) &= h(X \circ c, X \circ c) + h(dR_t \cdot \dot{\gamma}_0, X \circ c) \\ &= |X \circ c|_h^2 + h(dR_t \cdot \dot{\gamma}_0, dR_t \cdot X \circ \gamma_0) \\ &= |X \circ c|_h^2 + h(\dot{\gamma}_0, X \circ \gamma_0) > 0 \text{ } (\geq 0). \end{aligned}$$

□

Fix a point x_0 in the equator and let $0 < \phi_0 \leq \pi/2$ be determined as follows: the unique vector $v_{\phi_0} \in T_{x_0} S$ pointing to the northern hemisphere, satisfying $F(v_{\phi_0}) = 1$ and $h(v_{\phi_0} - X, X) = 0$ makes h -angle ϕ_0 with X . If for every $\phi \in [0, \phi_0]$ we denote by v_ϕ the unique vector not pointing south, satisfying $F(v_\phi) = 1$ and making h -angle ϕ with X , then $\phi < \phi_0 \Rightarrow h(v_\phi - X, X) > 0$. Analogously, we write c_ϕ for the geodesic of F satisfying $\dot{c}_\phi(0) = v_\phi$, and $\theta_\phi(t)$ for the unique lift of the function $\theta \circ c_\phi(t)$ to the universal covering $\mathbb{R}$ satisfying $\theta_\phi(0) = 0$, $\forall \phi \in [0, \phi_0]$. The lemma above provides the estimate $\dot{\theta}_\phi > 0$.

By (36), $c_{\phi_0}(t) = R_t \circ \gamma_+(t)$ where γ_+ is a geodesic of h heading north that leaves x_0 h -perpendicularly to the equator, and R_t is the flow of X . Thus c_{ϕ_0} passes through the north pole, and that is why $\theta_\phi(t)$ is defined only for $\phi < \phi_0$. Moreover, $1 = F(v_{\phi_0}) = |v_{\phi_0} - X|_h = |\dot{\gamma}_+(0)|_h$, see Remark 3.2.

Let $T_\phi > 0$ denote the first return time of the geodesic c_ϕ to the equator, and $P(\phi)$ be the point of return, which are well-defined smooth functions of $\phi \in (0, \phi_0]$. This is so since, by uniqueness of solutions of ODEs, the first hit of any geodesic c_ϕ , with $\phi \in (0, \phi_0]$, with the equator is transverse. We have that T_{ϕ_0} is equal to the h -length of the meridian T . To see that one needs to make use of the identity $F(v) = |v - X|_h$. For $\phi < \phi_0$ we have

$$P(\phi) = (R \cos \theta_\phi(T_\phi), R \sin \theta_\phi(T_\phi), 0).$$

Clearly the formula $c_{\phi_0}(t) = R_t \circ \gamma_+(t)$ implies

$$P(\phi_0) = (R \cos(\pi + T_{\phi_0} \eta), R \sin(\pi + T_{\phi_0} \eta), 0).$$

Thus $\text{dist}(\theta_\phi(T_\phi), \{\pi + T_{\phi_0} \eta + \{0, 2\pi, 4\pi, \dots\}\}) \rightarrow 0$ as $\phi \rightarrow \phi_0^-$. We used that $\dot{\theta}_\phi > 0$ for $\phi < \phi_0$. The curves c_ϕ converge in C_{loc}^1 to c_{ϕ_0} as $\phi \rightarrow \phi_0^-$, and c_{ϕ_0} does not self-intersect before hitting the equator since $T_{\phi_0} \eta < \pi$ by (43). Thus c_ϕ does not have a self-intersection before first hitting the equator when ϕ is close to ϕ_0 . This proves that $\theta_\phi(T_\phi) \rightarrow \pi + T_{\phi_0} \eta$ as $\phi \rightarrow \phi_0^-$.

The geodesic flow is the Reeb flow in the unit sphere bundle $SS^2 = F^{-1}(1)$ equipped with the contact form α_F discussed in § 2.3. The Jacobi vector field $J(t) = \partial_\phi|_{\phi=0} c_\phi(t)$ along $c_0(t)$ satisfies $g_{\dot{c}_0}(\dot{c}_0, J(t)) \equiv 0$ because it comes from a vertical variation and, as such, lies in the contact structure $\xi_F = \ker \alpha_F$. Thus $J(t) = f(t)\dot{c}_0^\perp(t)$, where f satisfies (27). Here g_v , $v \neq 0$, is the positive inner-product (15) on $T_{\pi(v)}S^2$, and $\dot{c}_0^\perp$ is the unique vector such that $\{\dot{c}_0, \dot{c}_0^\perp\}$ is a positively oriented $g_{\dot{c}_0}$ -orthonormal basis of $T_{c_0}S^2$. Since the flag curvatures along the equator are constant equal to 1, $f(t)$ is a (positive) multiple of $\sin t$ and, consequently, the first zero of J appears at time π . Thus

$$P(\phi) \rightarrow c_0(\pi) = (R \cos(\pi R^{-1} + \pi\eta), R \sin(\pi R^{-1} + \pi\eta), 0)$$

and $\theta_\phi(T_\phi) \rightarrow \pi R^{-1} + \pi\eta$ as $\phi \rightarrow 0^+$. Summarizing, we proved

$$(44) \quad \lim_{\phi \rightarrow \phi_0^-} \theta_\phi(T_\phi) = \pi + T_{\phi_0}\eta < 2\pi \quad (\text{see (43)})$$

and

$$(45) \quad \lim_{\phi \rightarrow 0^+} \theta_\phi(T_\phi) = \pi R^{-1} + \eta\pi = R^{-1}\pi(1 + \eta R) > \left(\frac{r}{r+1}\right)^{-1} \pi \frac{2r}{r+1} = 2\pi.$$

By continuity of $\phi \mapsto \theta_\phi(T_\phi)$, there exists $\phi^* \in (0, \phi_0)$ such that $\theta_{\phi^*}(T_{\phi^*}) = 2\pi$, and c_{ϕ^*} first returns to the equator exactly at x_0 . Moreover, c_{ϕ^*} does not self-intersect before hitting the equator since, otherwise, there would be some $t < T_{\phi^*}$ such that $\dot{\theta}_{\phi^*}(t) \leq 0$, a contradiction.

By the symmetry of F under the reflection Q with respect to the xy -plane,

$$\dot{c}_{\phi^*}(T_{\phi^*}) = dQ \cdot \dot{c}_{\phi^*}(0).$$

Here one has to make use of the Clairaut integral for the underlying Riemannian metric h . Thus, $c_\phi^* : [0, 2T_{\phi^*}] \rightarrow S^2$ is a smooth closed geodesic with precisely one transverse self-intersection. The flag curvatures lie between 1 and K_{max} , where K_{max} satisfies (40). We can normalize the curvature, after dilating the Finsler metric, in order to complete the proof of Theorem 1.5.

Remark 3.4. Given $r \geq 1$ and $0 < \delta < (r/(r+1))^2$ we can construct a surface of revolution $S_{\delta,r}$ as above to find an example of a δ -pinched Finsler metric on S^2 with reversibility equal to r , which is not dynamically convex. In fact, the double cover of the equator of $S_{\delta,r}$ corresponds to a contractible closed geodesic on its unit tangent bundle and has Conley-Zender index equal to 1. Therefore, the pinching condition on the flag curvatures given by Harris-Paternain in Theorem 1.8 that ensure dynamical convexity for α_F is sharp.

4. PROOF OF THEOREM 1.6

To prove the first assertion, observe that we can assume by contradiction the existence of convex hypersurfaces $S_k \subset \mathbb{R}^{2m}$, $k \geq 1$, converging to S^{2m-1} in $\mathbf{Conv}(2m)$ as $k \rightarrow \infty$, such that the Hamiltonian flow on S_k admits a closed orbit P_k with $\mathcal{A}(P_k) = \int_{P_k} \lambda_0 \rightarrow T_0 \in \mathbb{R}^+ \setminus \{\pi\}$. These corresponds to the existence of functions $f_k : S^{2m-1} \rightarrow (0, +\infty)$ converging to 1 in the C^2 -topology such that the Reeb flow $\Phi^k : \mathbb{R} \times S^{2m-1} \rightarrow S^{2m-1}$ associated to the contact form $\lambda_k := f_k \lambda_0|_{S^{2m-1}}$ admits a closed orbit, also denoted by P_k , with prime period T_k and satisfying $\mathcal{A}(P_k) = \int_{P_k} \lambda_k = T_k \rightarrow T_0 \in \mathbb{R}^+ \setminus \{\pi\}$. The Reeb vector fields R_0, R_k , $k \geq 1$, associated to $\lambda_0|_{S^{2m-1}}, \lambda_k$, $k \geq 1$, satisfy $R_k \rightarrow R_0$, as $k \rightarrow \infty$, in the C^1 -topology.

Since all the orbits of the Reeb flow Φ^0 associated to $\lambda_0|_{S^{2m-1}}$ are closed with prime period π , and since $T_k \rightarrow T_0$, we conclude from Arzelà-Ascoli theorem the existence of a simple closed orbit $P_0 \subset S^{2m-1}$ of Φ^0 and a subsequence of P_k , again denoted by P_k , such that P_k converges to a k_0 cover of P_0 , for some integer $k_0 > 1$ satisfying $T_0 = k_0\pi$. The following proposition due to Bangert [3] is crucial for completing the proof.

Proposition 4.1 (Bangert). *Let $\Phi : \mathbb{R} \times M \rightarrow M$, with M closed, be a C^1 flow and $p \in M$ be a periodic point of Φ with prime period T . Then for every $\epsilon > 0$ there exists a neighborhood $\mathcal{U}$ of Φ in the weak C^1 topology in $C^1(\mathbb{R} \times M, M)$ and a neighborhood U of p in M such that the following holds: if a flow $\hat{\Phi} \in \mathcal{U}$ has a periodic point $\hat{p} \in U$ with prime period $\hat{T}$ then either $\hat{T} > \epsilon^{-1}$ or there exists an integer $\hat{k} > 0$ such that $|\hat{T} - \hat{k}T| < \epsilon$ and the eigenvalues of the linear map*

$$(46) \quad D_2\Phi(T, p) : T_p M \rightarrow T_p M$$

which are $\hat{k}$ 'th roots of unity generate all the $\hat{k}$ 'th roots of unity.

Applying Proposition 4.1 to our situation we conclude that

$$(47) \quad D_2\Phi^0(\pi, p_0) = Id_{T_{p_0}S^{2m-1}} : T_{p_0}S^{2m-1} \rightarrow T_{p_0}S^{2m-1}, p_0 \in P_0,$$

admits an eigenvalue λ_0 which generates all the k_0 'th roots of unity. But this is a contradiction since $k_0 > 1$ and all eigenvalues of (47) are equal to 1.

To prove the statement made in the case $2m = 4$ about the linking numbers of short and long orbits we proceed indirectly and assume, by contradiction, the existence of $S_n \rightarrow S^3$ in $\mathbf{Conv}(4)$, $P_n, P'_n \in \mathcal{P}(S_n)$ satisfying $\mathcal{A}(P_n) \rightarrow \pi$, $\mathcal{A}(P'_n) \rightarrow +\infty$ and $\sup_n \text{link}(P'_n, P_n) = k_0 < \infty$.

We may view P_n and P'_n as closed Reeb orbits in S^3 of contact forms $f_n\lambda_0|_{S^3}$ with $f_n \rightarrow 1$ in the C^2 -topology. Let R_0 and R_n be the Reeb vector fields associated to $\lambda_0|_{S^3}$ and $f_n\lambda_0|_{S^3}$, respectively. Then $R_n \rightarrow R_0$ in the C^1 -topology. We denote by Φ_t^n and Φ_t^0 the flows of R_n, R_0 respectively. We can also assume, in view of the Arzelà-Ascoli theorem, that $P_n \rightarrow L_0$ in C^1 for some Hopf fiber $L_0 \subset S^3$.

The Hopf fiber L_0 corresponds to a π -periodic orbit of the flow Φ^0 . Identifying $\mathbb{R}^4 \simeq \mathbb{C}^2$, there is no loss of generality if we assume $L_0 = \{(e^{i\vartheta}, 0) \mid \vartheta \in \mathbb{R}\}$. Denote by $E : \mathbb{C}^2 \setminus \{0\} \rightarrow S^3$ the projection $E(x) = x/|x|$ and by Π_τ the plane $\{e^{i2\pi\tau}\} \times \mathbb{C}$. Then each $D_\tau := \overline{E(\Pi_\tau)}$ is an embedded disk, transverse to the vector field R_0 in $\mathring{D}_\tau$, satisfying $\partial D_\tau = L_1$, where L_1 is the Hopf fiber $L_1 = \{(0, e^{i\vartheta}) \mid \vartheta \in \mathbb{R}\}$. Moreover, D_τ is a global surface of section for Φ^0 and the (first) return map to D_τ is precisely the identity.

This open book decomposition induces a diffeomorphism

$$(48) \quad \begin{aligned} \Psi : S^3 \setminus L_1 &\rightarrow \mathbb{R}/\pi\mathbb{Z} \times \mathbb{C} \\ (z, w) &\mapsto \left(\frac{\arg z}{2}, \frac{w}{\sqrt{1-|w|^2}} \right) \end{aligned}$$

satisfying $\Psi(\mathring{D}_\tau) = \{\tau\} \times \mathbb{C}$. The flow $\hat{\Phi}_t^0 := \Psi_*\Phi_t^0 = \Psi \circ \Phi_t^0 \circ \Psi^{-1}$ is obtained by integrating the vector field $\hat{R}_0 := \Psi_*R_0$ and is given by

$$(49) \quad \hat{\Phi}_t^0(\tau_0, \zeta) = (\tau_0 + t, e^{i2t}\zeta).$$

Note that L_0 is mapped precisely onto $\mathbb{R}/\pi\mathbb{Z} \times 0$ since $\Psi(e^{i2\vartheta}, 0) = (\vartheta, 0)$.

Since $P_n \rightarrow L_0$ in C^1 , we find diffeomorphisms F_n of $\mathbb{R}/\pi\mathbb{Z} \times \mathbb{C}$ satisfying: $\text{supp}(F_n) \subset \mathbb{R}/\pi\mathbb{Z} \times B_1(0)$, $F_n \rightarrow \text{id}$ in C^1 and $F_n \circ \Psi(P_n) = \mathbb{R}/\pi\mathbb{Z} \times 0$. This follows from an application of Lemma 4.2.

Lemma 4.2. *Let M be a C^k -manifold and $Z \hookrightarrow M$ be a closed C^k -submanifold. Suppose U is any open neighborhood of Z and Z_n are C^k -submanifolds converging to Z in the C^k -topology as $n \rightarrow \infty$. Then there exist diffeomorphisms φ_n satisfying $\varphi_n(Z_n) = Z$, $\text{supp}(\varphi_n) \subset U$ and $\varphi_n \rightarrow \text{id}_M$ in the C^k -topology.*

Proof. By considering a tubular neighborhood of Z in M one sees that there is no loss of generality if we assume M is C^k -vector bundle, Z is the zero section, U is a neighborhood of the zero section and the Z_n are graphs of sections s_n converging to the zero section in the C^k -topology. Let $f : M \rightarrow [0, 1]$ be a fixed smooth function with support compactly contained in U that is identically equal to 1 in a neighborhood of the zero section containing all Z_n . If π is the projection of M onto its base, we define the diffeomorphism $\varphi_n : M \rightarrow M$ by $\varphi_n(v) := v - f(v)s_n(\pi(v))$. It is easy to check that φ_n satisfies all requirements, when n is sufficiently large. $\square$

If we set $\Psi_n := F_n \circ \Psi$, $\hat{R}_n := (\Psi_n)_* R_n$ and denote by $\hat{\Phi}^n$ the flow of $\hat{R}_n$ then the maximal domain of definition of $\hat{\Phi}^n$ is an exhausting sequence of open subsets of $\mathbb{R} \times \mathbb{R}/\pi\mathbb{Z} \times \mathbb{C}$ and $\hat{\Phi}^n \rightarrow \hat{\Phi}^0$ in the C^1 -topology on compact sets of $\mathbb{R} \times \mathbb{R}/\pi\mathbb{Z} \times \mathbb{C}$.

Consider $D_0 := \overline{\Psi^{-1}(\mathbb{R}/\pi\mathbb{Z} \times [0, +\infty))}$. This set is a smooth embedded disk satisfying $\partial D_0 = L_0$ which is transverse to R_0 at $\mathring{D}_0 = D_0 \setminus L_0$. In fact, $\mathring{D}_0$ coincides with $E(\mathbb{C} \times \{1\})$. Now we consider disks

$$(50) \quad D_n := \overline{\Psi_n^{-1}(\mathbb{R}/\pi\mathbb{Z} \times [0, +\infty))}.$$

Since the support of F_n is contained in $\mathbb{R}/\pi\mathbb{Z} \times B_1(0)$ the disk D_n coincides with D_0 on $S^3 \setminus \Psi^{-1}(\mathbb{R}/\pi\mathbb{Z} \times B_1(0))$. Moreover, $\partial D_n = P_n$.

We claim D_n is transverse to R_n at the points of $D_n \setminus P_n$. Using Taylor's formula and comparing with $\hat{R}_0$ we obtain

$$(51) \quad \begin{aligned} \hat{R}_n(\tau, z) &= \hat{R}_n(\tau, 0) + D_2 \hat{R}_0(\tau, 0) \cdot z \\ &+ \left[D_2 \hat{R}_n(\tau, 0) - D_2 \hat{R}_0(\tau, 0) + \int_0^1 D_2 \hat{R}_n(\tau, \lambda z) - D_2 \hat{R}_n(\tau, 0) d\lambda \right] \cdot z \\ &= \hat{R}_n(\tau, 0) + D_2 \hat{R}_0(\tau, 0) \cdot z + \epsilon_n(\tau, z) \cdot z \end{aligned}$$

with

$$\sup\{|\epsilon_n(\tau, z)| : (\tau, z) \in \mathbb{R}/\pi\mathbb{Z} \times B_1(0)\} \rightarrow 0$$

as $n \rightarrow \infty$. In these coordinates the vector $(0, 0, 1)$ is normal to the strip $\mathbb{R}/\pi\mathbb{Z} \times [0, +\infty)$. Note that

$$(52) \quad D_2 \hat{R}_0(\tau, 0) = \begin{bmatrix} 0 & 0 \\ 0 & -2 \\ 2 & 0 \end{bmatrix}$$

so for every $\eta > 0$ we find $r_0 > 0$ and $n_0 > 0$ such that

$$(53) \quad \left\langle \hat{R}_n(\tau, r), (0, 0, 1) \right\rangle \geq (2 - \eta)r$$

for every $n \geq n_0$ and $r < r_0$. This shows that $\hat{R}_n$ is transverse to $\mathbb{R}/\pi\mathbb{Z} \times (0, r_0)$ if n is large since we can choose $\eta < 2$, which amounts to say that $\Psi_n^{-1}(\mathbb{R}/\pi\mathbb{Z} \times (0, r_0))$

is transverse to R_n . Now observe that $\overline{\Psi_n^{-1}(\mathbb{R}/\pi\mathbb{Z} \times [r_0, +\infty))}$ is converging in the C^1 -topology to the disk $\overline{\Psi^{-1}(\mathbb{R}/\pi\mathbb{Z} \times [r_0, +\infty))}$, which is transverse to R_0 . So $\Psi_n^{-1}(\mathbb{R}/\pi\mathbb{Z} \times [r_0, +\infty))$ will also be transverse to R_n if n is large, proving our claim.

Remark 4.3. Fix n and suppose $\gamma : [a, b] \rightarrow S^3 \setminus (P_n \cup L_1)$ is a closed curve. Define $s \in [a, b] \mapsto \zeta(s) \in \mathbb{C} \setminus \{0\}$ by $\Psi_n \circ \gamma(s) = (*, \zeta(s))$. Then it is not hard to check that

$$(54) \quad \text{link}(\gamma, P_n) = \frac{\vartheta(b) - \vartheta(a)}{2\pi}$$

where $\vartheta : [a, b] \rightarrow \mathbb{R}$ is any continuous lift of the argument of $\zeta(s)$.

We split the remaining arguments in two cases.

Case 1: $\forall n \exists y_n \in P'_n$ satisfying $\text{dist}(y_n, L_1) \rightarrow 0$.

Since L_1 is a closed orbit of R_0 , we can assume the existence of $z_1 \in L_1 \cap D_0$ and $y_n \in P'_n$ such that $y_n \rightarrow z_1$ as $n \rightarrow \infty$. Let $V \subset D_0$ be a small neighborhood of z_1 . Note that by the construction above, $V \subset D_n$ for all n large. Now since $R_n \rightarrow R_0$ and $\Phi_\pi^0(z_1) = z_1$, given any integer $k > 0$ and any real number $\epsilon > 0$ we can find $U_k \subset V$ neighborhood of z_1 , $n_0(k, \epsilon) > 0$, both depending on k and ϵ , such that for all $n > n_0(k, \epsilon)$ and $z \in U_{k, \epsilon}$, the solution $\{\Phi_t^n(z), t \in (0, (k + 1/2)\pi]\}$, intersects V transversely and positively at least k times, and these intersections correspond to points $t \in [j\pi - \epsilon, j\pi + \epsilon]$ for each $j \in \{1, \dots, k\}$. Now since $\mathcal{A}(P'_n) \rightarrow \infty$ as $n \rightarrow \infty$, it follows that $\text{link}(P_n, P'_n) = \#\{P'_n \cap D_n\} \rightarrow \infty$ as $n \rightarrow \infty$ which is a contradiction. In this last assertion we strongly used that each D_n is a (positively) transverse disk to R_n , when n is large, so P'_n never intersects D_n negatively.

Case 2: $\inf\{\text{dist}(y, L_1) \mid y \in P'_n\} \geq \delta > 0$.

By our hypotheses $\exists \rho > 0$ such that $\Psi(P'_n) \subset \mathbb{R}/\pi\mathbb{Z} \times B_\rho(0)$ when n is large enough. Moreover, $\hat{P}'_n := \Psi_n(P'_n)$ is a T'_n -periodic orbit of the flow $\hat{\Phi}_t^n$ completely contained in $\mathbb{R}/\pi\mathbb{Z} \times B_\rho(0)$, and $T'_n \rightarrow \infty$.

For each n fix a point $y_n \in \hat{P}'_n \cap (0 \times B_\rho(0))$. Define $\gamma_n(t) := \hat{\Phi}_t^n(y_n) = (\tau_n(t), \zeta_n(t)) \in \mathbb{R}/\pi\mathbb{Z} \times B_\rho(0)$. If $\tilde{\tau}_n(t)$ is a continuous lift of $\tau_n(t)$ and $\vartheta_n(t)$ is a continuous lift of the argument of $\zeta_n(t)$, then we consider

$$N_n := \frac{\tilde{\tau}_n(T'_n) - \tilde{\tau}_n(0)}{\pi} \in \mathbb{Z}^+ \quad \text{and} \quad l_n := \frac{\vartheta_n(T'_n) - \vartheta_n(0)}{2\pi} \in \mathbb{Z}.$$

By (54) we have $l_n = \text{link}(P'_n, P_n)$.

The (first) return time with respect to the flow $\hat{\Phi}_t^n$ for points of $0 \times B_\rho(0)$ to return to $0 \times \mathbb{C}$ is a function converging uniformly to the constant π on $0 \times B_\rho(0)$ since $\hat{\Phi}^n \rightarrow \hat{\Phi}^0$ as above. Since $\hat{R}_n \rightarrow \hat{R}_0$ in C_{loc}^1 , $\hat{R}_n$ is transverse to the disks $\tau \times B_\rho(0)$, $\tau \in \mathbb{R}/\pi\mathbb{Z}$, when n is large enough. Consequently we can divide, for each n large enough, the interval $[0, T'_n]$ in precisely N_n intervals $\{I_1^n, \dots, I_{N_n}^n\}$ of lengths uniformly close to π corresponding to points where P'_n intersects $0 \times B_\rho(0)$. This implies that $N_n \rightarrow \infty$ as $n \rightarrow \infty$.

We need to estimate $\dot{\vartheta}_n$. Let us write $\hat{R}_n = (a_n, Y_n = (u_n, v_n))$. If $\zeta_n(t) = x_n(t) + iy_n(t)$ then

$$\begin{aligned}\dot{\vartheta}_n &= \frac{-y_n \dot{x}_n + x_n \dot{y}_n}{x_n^2 + y_n^2} = \frac{\langle \zeta_n, -J_0 \cdot \dot{\zeta}_n \rangle}{|\zeta_n|^2} \\ &= \frac{\langle \zeta_n, -J_0 \cdot (Y_n \circ \gamma_n) \rangle}{|\zeta_n|^2} \\ &= \left\langle \frac{\zeta_n}{|\zeta_n|}, -J_0 \cdot [DY_n(\tau_n(t), 0) + \epsilon_n] \cdot \frac{\zeta_n}{|\zeta_n|} \right\rangle\end{aligned}$$

where $J_0 = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$ and

$$\epsilon_n(t) = \int_0^1 [DY_n(\tau_n(t), \lambda \zeta_n(t)) - DY_n(\tau_n(t), 0)] d\lambda.$$

Now, since $\hat{R}_n$ converges C^1 to $\hat{R}_0(\tau, z) = (1, i2z)$, uniformly on $\mathbb{R}/\pi\mathbb{Z} \times \overline{B_\rho(0)}$, we conclude that, as $n \rightarrow \infty$, ϵ_n converges uniformly in $t \in [0, T'_n]$ to the zero matrix and $DY_n(\tau_n(t), 0)$ converges uniformly in $t \in [0, T'_n]$ to the matrix

$$\begin{bmatrix} 0 & -2 \\ 2 & 0 \end{bmatrix}$$

as was computed in (52). Consequently, one estimates $\dot{\vartheta}_n \geq 3/2$, uniformly in $t \in [0, T'_n]$. Thus ϑ_n is strictly increasing, and increases at least $3\pi/2$ on each I_j^n , for every j and every n sufficiently large. Thus $l_n \rightarrow +\infty$, and this contradiction concludes Case 2.

APPENDIX A. LEMMAS FROM FINSLER GEOMETRY

A.1. Proof of Lemma 2.1. For a fixed j we have, in natural coordinates, $(\partial_{x^j})_h = \partial_{x^j} - \Gamma_j^k \partial_{y^k}$. Thus, since S is horizontal, we obtain

$$\begin{aligned}[P_H(S), P_H((\partial_{x^j})_h)] &= [S, (\partial_{x^j})_h] = [y^i \partial_{x^i} - 2G^i \partial_{y^i}, \partial_{x^j} - \Gamma_j^k \partial_{y^k}] \\ &= -2[G^i \partial_{y^i}, \partial_{x^j}] - [y^i \partial_{x^i}, \Gamma_j^k \partial_{y^k}] + 2[G^i \partial_{y^i}, \Gamma_j^k \partial_{y^k}] \\ &= 2(\partial_{x^j} G^i) \partial_{y^i} - y^i (\partial_{x^i} \Gamma_j^k) \partial_{y^k} - \Gamma_j^k [y^i \partial_{x^i}, \partial_{y^k}] + 2G^i (\partial_{y^i} \Gamma_j^k) \partial_{y^k} + 2\Gamma_j^k [G^i \partial_{y^i}, \partial_{y^k}] \\ &= 2(\partial_{x^j} G^i) \partial_{y^i} - y^i (\partial_{x^i} \Gamma_j^k) \partial_{y^k} + \Gamma_j^i \partial_{x^i} + 2G^i (\partial_{y^i} \Gamma_j^k) \partial_{y^k} - 2\Gamma_j^k (\partial_{y^k} G^i) \partial_{y^i} \\ &= \{2(\partial_{x^j} G^i) - y^k (\partial_{x^k} \Gamma_j^i) + 2G^k (\partial_{y^k} \Gamma_j^i) - 2\Gamma_j^k (\partial_{y^k} G^i)\} \partial_{y^i} + \Gamma_j^i \partial_{x^i}\end{aligned}$$

and consequently

$$\begin{aligned}(55) \quad R(S, (\partial_{x^j})_h) &= P_V([P_H(S), P_H((\partial_{x^j})_h)]) \\ &= \{2(\partial_{x^j} G^i) - y^k (\partial_{x^k} \Gamma_j^i) + 2G^k \Gamma_{jk}^i - 2\Gamma_j^k \Gamma_k^i + \Gamma_j^i \Gamma_k^k\} \partial_{y^i}\end{aligned}$$

Here we used $P_V(\partial_{x^i}) = \Gamma_i^k \partial_{y^k}$. On the other hand,

$$\begin{aligned}(56) \quad \nabla_{[S, (\partial_{x^j})_h]} C &= \nabla_{[S, (\partial_{x^j})_h]} (y^l \partial_{y^l}) = (dy^l \cdot [S, (\partial_{x^j})_h]) \partial_{y^l} + y^l \nabla_{[S, (\partial_{x^j})_h]} \partial_{y^l} \\ &= (dy^l \cdot [S, (\partial_{x^j})_h]) \partial_{y^l} + y^l \Gamma_j^i \nabla_{\partial_{x^i}} \partial_{y^l} = (dy^l \cdot [S, (\partial_{x^j})_h]) \partial_{y^l} + y^l \Gamma_j^i \Gamma_{il}^k \partial_{y^k} \\ &= \{2(\partial_{x^j} G^l) - y^k (\partial_{x^k} \Gamma_j^l) + 2G^k \Gamma_{jk}^l - 2\Gamma_j^k \Gamma_k^l + y^k \Gamma_j^i \Gamma_{ik}^l\} \partial_{y^l}\end{aligned}$$

Using the identity $y^k \Gamma_{ik}^l = \Gamma_i^l$ one sees that (55) equals (56). Since $\nabla_S C = \nabla_{(\partial_{x^j})_h} C = 0$ one computes at the base point v :

$$\tilde{R}(v_h, (\partial_{x^j})_h) i_v(v) = \tilde{R}(S, (\partial_{x^j})_h) C = -\nabla_{[S, (\partial_{x^j})_h]} C = -R(S, (\partial_{x^j})_h).$$

Here we used that the horizontal lift of v to $T_v TM$ is S , that $i_v(v) = C$ and, in the last equality, that (55) equals (56). The conclusion follows because i_v is an isomorphism.

A.2. Proof of Lemma 2.2. To prove (22) take natural coordinates and write

$$\text{LHS} = ((\partial_{x^l} g_{ij}) \dot{x}^l - 2(\partial_{y^l} g_{ij}) G^l - g_{lj} \Gamma_i^l - g_{il} \Gamma_j^l) V^i W^j + \text{RHS}.$$

Since V and W are arbitrary we get

$$(\partial_{x^l} g_{ij}) \dot{x}^l - 2(\partial_{y^l} g_{ij}) G^l - g_{lj} \Gamma_i^l - g_{il} \Gamma_j^l = 0, \quad \forall i, j.$$

Choosing a symmetric lift ∇ of Γ_S , with local coefficients Γ_{ij}^l , the above expression becomes $((\partial_{x^k} g_{ij}) - A_{ijl}(\partial_{y^k} G^l) - g_{lj} \Gamma_{ik}^l - g_{il} \Gamma_{jk}^l) \dot{x}^k = 0$, for every i, j, k , where $A_{ijl} = \frac{1}{2} \partial_{y^l} g_{ij}$ are the components of the Cartan tensor (20). Thus, if the symmetric lift satisfies

$$(57) \quad (\partial_{x^k} g_{ij}) - 2A_{ijl}(\partial_{y^k} G^l) - g_{lj} \Gamma_{ik}^l - g_{il} \Gamma_{jk}^l = 0, \quad \forall i, j, k$$

the claim follows. However, this may not be satisfied by an arbitrary ∇ . In fact, we can permute the (i, j, k) above to get terms corresponding to (j, k, i) and (k, i, j) . Adding the (i, j, k) -term to the (j, k, i) -term, subtracting the (k, i, j) -term and using $\Gamma_m^l = \partial_{y^m} G^l$, one obtains

$$2g_{jl} \Gamma_{ik}^l = \partial_{x^k} g_{ij} - \partial_{x^j} g_{ki} + \partial_{x^i} g_{jk} - 2\{A_{ijl} \Gamma_k^l + A_{jkl} \Gamma_i^l - A_{kil} \Gamma_j^l\}.$$

Multiplying by $\frac{1}{2} g^{mj}$ and summing in j we get equations (21) for the coefficients of the Chern connection, that is, if we use the Chern connection as the symmetric lift, the desired conclusion holds. However, equation (22) does not depend on this choice, which implies that we could have chosen any symmetric lift to carry on our calculations.

A.3. Proof of Lemma 2.3. Consider, in natural coordinates $x = (x^1, \dots, x^n)$, $y = (y^1, \dots, y^n)$, the ODE $(\dot{x}, \dot{y}) = (y, -2G)$ associated to the geodesic spray, where $G = (G^1, \dots, G^n)$ are the spray coefficients. Linearizing we get

$$(58) \quad \begin{pmatrix} \delta x \\ \delta y \end{pmatrix} = \begin{pmatrix} 0 & I \\ -2D_x G & -2D_y G \end{pmatrix} \begin{pmatrix} \delta x \\ \delta y \end{pmatrix}$$

where $\delta x = (\delta x^1, \dots, \delta x^n)$ and $\delta y = (\delta y^1, \dots, \delta y^n)$ are the fiber coordinates on TTM . Here $D_x G$ and $D_y G$ are $n \times n$ matrices with entries $\partial_{x^i} G^k$ and $\partial_{y^i} G^k$, respectively, evaluated at $(x^1 = x^1 \circ \gamma(t), \dots, y^1(t) = \dot{x}^1, \dots)$, where $\gamma(t)$ is some geodesic.

We need to rewrite (58) in terms of invariantly defined objects. Consider the vector field $J(t) = \delta x^i \partial_{x^i}$ along γ . Then $\delta \dot{x}^k = (\frac{D_{\gamma} J}{dt})^k - \Gamma_l^k \delta x^l$, where $\frac{D_{\gamma} J}{dt} =$

$(\frac{D_\gamma J}{dt})^k \partial_{x^k}$, and

$$\begin{aligned}
 \delta \ddot{x}^k &= \frac{d}{dt} \left[\left(\frac{D_\gamma J}{dt} \right)^k - \Gamma_l^k \delta x^l \right] = \left(\frac{D_\gamma^2 J}{dt^2} \right)^k - \Gamma_l^k \left(\frac{D_\gamma J}{dt} \right)^l - \frac{d}{dt} (\Gamma_l^k \delta x^l) \\
 &= \left(\frac{D_\gamma^2 J}{dt^2} \right)^k - 2\Gamma_l^k \dot{\delta x}^l - (\partial_{x^i} \Gamma_l^k) \dot{x}^i \delta x^l + 2(\partial_{y^i} \Gamma_l^k) G^i \delta x^l - \Gamma_i^k \Gamma_l^i \delta x^l
 \end{aligned}
 \tag{59}$$

where we used $\dot{y}^i + 2G^i = 0$. Adding $2(\partial_{x^i} G^k) \delta x^i + 2(\partial_{y^i} G^k) \dot{\delta x}^i$ to (59) we get 0, in view of (58), and

$$\begin{aligned}
 0 &= \left(\frac{D_\gamma^2 J}{dt^2} \right)^k - 2\Gamma_l^k \dot{\delta x}^l - (\partial_{x^i} \Gamma_l^k) \dot{x}^i \delta x^l + 2(\partial_{y^i} \Gamma_l^k) G^i \delta x^l \\
 &\quad - \Gamma_i^k \Gamma_l^i \delta x^l + 2(\partial_{x^i} G^k) \delta x^i + 2(\partial_{y^i} G^k) \dot{\delta x}^i \\
 &= \left(\frac{D_\gamma^2 J}{dt^2} \right)^k + \{ 2(\partial_{x^i} G^k) - (\partial_{x^i} \Gamma_l^k) \dot{x}^i + 2G^i (\partial_{y^i} \Gamma_l^k) - \Gamma_l^i \Gamma_i^k \} \delta x^l.
 \end{aligned}
 \tag{60}$$

Now observe that $R^\dot{\gamma}(J) = \delta x^l R^\dot{\gamma}(\partial_{x^l}) = \delta x^l i_{\dot{\gamma}}^{-1}(-\nabla_{[S, (\partial_{x^l})_h]} C)$ where the vectors $\nabla_{[S, (\partial_{x^l})_h]} C$ were computed in (56). Plugging into (60) we get equations

$$0 = \left(\frac{D_\gamma^2 J}{dt^2} \right)^k - R^\dot{\gamma}(J)^k, \quad \forall k = 1 \dots n.$$

REFERENCES

- [1] S. Angenent. *Curve Shortening and the topology of closed geodesics on surfaces*. Ann. of Math. (2) **162** (2005), 1185–1239.
- [2] V. I. Arnol'd. *Topological Invariants of Plane Curves and Caustics*. Dean Jacqueline B. Lewis Memorial Lectures presented at Rutgers University, New Brunswick, New Jersey. University Lecture Series, **5**. American Mathematical Society, Providence, RI, 1994. viii+60 pp.
- [3] V. Bangert. *On the lengths of closed geodesics on almost round spheres*. Math. Z. **191** (1986), no. 4, 549–558.
- [4] W. Ballmann. *On the length of closed geodesics on convex surfaces*. Invent. Math. **71** (1983), 593–597.
- [5] W. Ballmann, G. Thorbergsson and W. Ziller. *Closed geodesics on positively curved manifolds*. Ann. of Math. **116** (1982), 213–247.
- [6] W. Ballmann, G. Thorbergsson and W. Ziller. *Existence of closed geodesics on positively curved manifolds*. J. Differential Geom. **18** (1983), 221–252.
- [7] W. Ballmann, G. Thorbergsson and W. Ziller. *Some existence theorems for closed geodesics*. Comment. Math. Helvetici **58** (1983), 416–432.
- [8] D. Bao and C. Robles. *Ricci and flag curvatures in Finsler geometry*. A sampler of Riemann-Finsler geometry, 197–259, Math. Sci. Res. Inst. Publ., **50**, Cambridge Univ. Press, Cambridge, 2004.
- [9] G. D. Birkhoff. *Dynamical systems.*, (1927), AMS.
- [10] G. Contreas and F. Oliveira. *C^2 -densely the 2-sphere has an elliptic closed geodesic*. Ergod. Th. & Dynam. Sys. **24** (2004), 1395–1423.
- [11] J. Grifone. *Structure presque-tangente et connexions. I*. Ann. Inst. Fourier (Grenoble) **22** (1972), no. 1, 287–334.
- [12] M. Gromov. *Pseudoholomorphic curves in symplectic manifolds*. Invent. Math. **82** (1985), 307–347.
- [13] A. Harris and G. Paternain. *Dynamically convex Finsler metrics and J -holomorphic embedding of asymptotic cylinders*. Ann. Global Anal. Geom. **34** (2008), no. 2, 115–134.

- [14] H. Hofer. *Pseudoholomorphic curves in symplectisations with application to the Weinstein conjecture in dimension three*. Invent. Math. **114** (1993), 515–563.
- [15] H. Hofer, K. Wysocki and E. Zehnder. *Properties of pseudoholomorphic curves in symplectisations I: Asymptotics*. Ann. Inst. H. Poincaré Anal. Non Linéaire **13** (1996), 337–379.
- [16] H. Hofer, K. Wysocki and E. Zehnder. *A characterization of the tight three sphere*. Duke Math. J. **81** (1995), no. 1, 159–226.
- [17] H. Hofer, K. Wysocki and E. Zehnder. *A characterization of the tight three sphere II*. Commun. Pure Appl. Anal. **55** (1999), no. 9, 1139–1177.
- [18] H. Hofer, K. Wysocki and E. Zehnder. *The dynamics of strictly convex energy surfaces in $\mathbb{R}^4$* . Ann. of Math. **148** (1998), 197–289.
- [19] H. Hofer, K. Wysocki and E. Zehnder. *Finite energy foliations of tight three-spheres and Hamiltonian dynamics*. Ann. Math **157** (2003), 125–255.
- [20] U. Hryniewicz. *Fast finite-energy planes in symplectizations and applications*. To appear in Trans. Amer. Math. Soc. (arXiv:0812.4076).
- [21] A. Katok. *Ergodic properties of degenerate integrable Hamiltonian systems*. Izv. Akad. Nauk SSSR. **37** (1973) (Russian), 535–571.
- [22] W. Klingenberg. *Der Indexsatz für geschlossene Geodätsche*. Math. Z. **139** (1974), 231–256.
- [23] H. Poincaré. *Sur les lignes géodésiques des surfaces convexes*. Trans. Amer. Math. Soc. **6** (1905), 237–274.
- [24] H.-B. Rademacher. *A sphere theorem for non-reversible Finsler metrics*. Math. Ann. **328** (2004), 373–387.
- [25] H.-B. Rademacher. *Nonreversible Finsler metrics of positive flag curvature*. A sampler of Riemann-Finsler geometry, 261–302, Math. Sci. Res. Inst. Publ. **50**, Cambridge Univ. Press, Cambridge, 2004.
- [26] H.-B. Rademacher. *The length of a shortest geodesic loop*. Comptes Rendus Mathematique, Volume **346**, 13–14 (2008), 763–765.
- [27] G. Thorbergsson. *Non-hyperbolic closed geodesics*. Math. Scand. **44** (1979), 135–148.
- [28] H. Vitória. *A geometria de curvas fanning e de suas reduções simpléticas*. Ph.D. Thesis, Unicamp, Campinas, 2010.

(Umberto L. Hryniewicz) INSTITUTE FOR ADVANCED STUDY/DEPARTAMENTO DE MATEMÁTICA APLICADA – UNIVERSIDADE FEDERAL DO RIO DE JANEIRO

E-mail address: umbertolh@math.ias.edu/umberto@labma.ufrj.br

(Pedro A. S. Salomão) DEPARTAMENTO DE MATEMÁTICA, INSTITUTO DE MATEMÁTICA E ESTATÍSTICA – UNIVERSIDADE DE SÃO PAULO

E-mail address: psalomao@ime.usp.br